Purpose and procedure

The general purpose of *Evidence-Based Nursing* is to select from the health-related literature those articles reporting studies and reviews that warrant immediate attention by nurses attempting to keep pace with important advances in their profession. These articles are summarised in "value added" abstracts and commented on by nurses. The specific purposes of *Evidence-Based Nursing* are

- ▶ to identify, using predefined criteria, the best quantitative and qualitative original and review articles on the meaning, cause, course, assessment, prevention, treatment, or economics of health problems managed by nurses and on quality improvement
- ▶ to summarise this literature in the form of "structured abstracts" that describe the question, methods, results, and evidence-based conclusions of studies in a reproducible and accurate fashion
- ▶ to provide brief, highly expert comment on the context of each article, its methods, and the clinical applications that its findings warrant
- ▶ to disseminate the summaries in a timely fashion to nurses

The RCN Publishing Company Limited and the BMJ Publishing Group publish *Evidence-Based Nursing* under the editorship of Dr Donna Ciliska at McMaster University in Canada, Dr Andrew Jull at the University of Auckland in New Zealand, and Dr Carl Thompson at the University of York in the UK. The Health Information Research Unit (HIRU) of the Department of Clinical Epidemiology and Biostatistics at McMaster University hosts the editorial office for the production of the abstracts and commissioning of commentaries. Dr Brian Haynes acts as coordinating editor to ensure that methods and procedures are consistent with other evidence-based journals prepared by HIRU.

CRITERIA FOR SELECTION AND REVIEW OF ARTICLES FOR ABSTRACTING

All articles in a journal issue are considered for abstracting if they meet the following basic and category-specific criteria:

Basic criteria

- original or review articles
- ▶ in English
- quantitative and qualitative studies
- ▶ about topics that are important to the clinical practice of nurses in any setting
- lacktriangle analysis of data consistent with the study question

Category-specific criteria (quantitative studies) Studies of prevention or treatment must also include:

- ▶ random allocation of participants to comparison groups
- follow-up (end-point assessment) of ≥80% of those entering the investigation
- ▶ outcome measures of known or probable clinical importance

Studies of assessment (screening or diagnosis) must also include:

- ▶ a spectrum of participants, some (but not all) of whom have the condition of interest
- an objective diagnostic (gold) standard (eg, central venous pressure) or current clinical standard for diagnosis (eg, sphygmomanometer reading for hypertension), preferably

- with documentation of reproducible criteria for subjectively interpreted diagnostic standard (ie, report of statistically significant measure of agreement beyond chance among observers)
- ► interpretation of the diagnostic standard without knowledge of test result
- interpretation of the test without knowledge of diagnostic standard result

Studies of prognosis must also include:

- an inception cohort (first onset or assembled at a uniform point in the development of a condition or disease) of individuals, all initially free of the outcome of interest
- ► follow-up of ≥80% of participants until a major study endpoint occurs or the study ends

Studies of causation must also include:

- ▶ observations of the relation between modifiable exposures and putative clinical outcomes
- ▶ prospective data collection with clearly identified comparison group(s) for those at risk of, or having, the outcome of interest (ie, randomised controlled trials, quasi-randomised controlled trials, non-randomised controlled trials, cohort studies with case by case matching or statistical adjustment to create comparable groups, or nested case—control studies)
- blinding (masking) of observers of outcome to exposure; criterion assumed to be met if outcome is objective (eg, allcause mortality)

Studies of quality improvement or continuing education must also include:

- random allocation of participants or units to comparison groups
- ► follow-up of ≥80% of participants
- outcome measures of known or probable clinical importance

Studies of the economics of healthcare interventions must include:

- ▶ an economic question comparing alternative courses of action
- ► comparison of alternative diagnostic or therapeutic services or quality improvement activities on the basis of both the outcomes produced (effectiveness) and resources consumed (costs)
- evidence of effectiveness from a study (or studies) of real (not hypothetical) patients, which meet(s) the criteria for treatment, assessment, quality improvement, or a systematic review article
- results presented in terms of the incremental or additional costs and outcomes of one intervention over another
- ► a sensitivity analysis if there is uncertainty in the estimates or imprecision in the measurement

Clinical prediction guides must also include:

- generation of the guide in ≥1 set of patients (training or derivation set)
- validation of the guide in an independent set of patients (test or validation set)

EBN October 2009 Vol 12 No 4

Systematic review articles must also include:

- a clear statement of the clinical topic being reviewed
- a clear description of the sources and methods for identifying articles
- specification of the inclusion and exclusion criteria for selecting articles for detailed review
- ≥1 article in the review that meets the above noted criteria for treatment, assessment, prognosis, causation, quality improvement, economics, or clinical prediction guides

Qualitative studies

- content reflects the phenomenon of interest from the perspective of people experiencing it
- data collection methods are appropriate for qualitative data
- analyses are appropriate for qualitative data

These criteria are subject to modification if, for example, it becomes feasible to apply higher standards that increase the validity and applicability of studies for clinical practice. The objective of Evidence-Based Nursing is to abstract only the very best literature, consistent with a reasonable number of articles "making it through the filter."

Articles meeting the criteria set out above are abstracted according to the procedure for more informative abstracts.¹ with the following modifications: abstracts are approximately 400 words in length; and each abstract is reviewed by an expert in the content area covered by the article. This expert writes a commentary in which she or he compares the study findings to previous research findings, identifies any important methodological problems that affect interpretation of the study results, and offers recommendations for clinical application. The author of the original article is given an opportunity to review the abstract and commentary before publication.

On an quarterly basis, we will publish to the Evidence-Based Nursing website (www.evidencebasednursing.com) a selected list of articles that passed all criteria but were not abstracted because, in the judgment of the editors, their findings were less applicable to general nursing practice, the topic was of interest to only a select group of nurse specialists, or the topic was recently addressed in another abstract.

Haynes RB, Mulrow CD, Huth EJ, et al. More informative abstracts revisited. Ann Intern Med 1990;113:69-76.

Journals currently reviewed*

Arthritis Rheum: Arthritis Acad Emerg Med Care Res Acta Obstet Gynecol Scand

Acta Orthop Age Ageing

Aliment Pharmacol Ther

Am J Cardiol Am J Clin Nutr Am J Epidemiol Am J Gastroenterol Am J Kidnev Dis

Am J Med Am J Obstet Gynecol Am J Occup Ther

Am J Psychiatry Am J Public Health Am J Respir Crit Care Med

Am J Sports Med **Anesth Analg** Anesthesiology

Ann Allerg Asthma Immunol Ann Emerg Med

Ann Fam Med Ann Intern Med Ann Neurol Ann Rheum Dis Ann Surg

ANS Adv Nurs Sci Appl Nurs Res Arch Dermatol Arch Dis Child

Arch Dis Child Fetal Neonatal Ed Arch Gen Psychiatry Arch Intern Med

Arch Neurol

Arch Pediatr Adolesc Med Arch Phys Med Rehabil

Arch Surg Arthritis Rheum

Arthroscopy Aust J Physiother

Birth **BJOG** BM.I Br J Gen Pract

Br J Psychiatry Br J Surg Can J Anaesth

Can J Gastroenterol Can J Nurs Res

Canadian Agency for Drugs and Technologies in Health (CADTH) Cancer Nurs

Cephalalgia Chest Circulation Clin Chem

Clin J Am Soc Nephrol

Clin J Pain Clin Orthop Rel Res

Clin Rehab

CMAJ

Cochrane Database Syst Rev

Crit Care Med Diabet Med Diabetes Care Diabetes Obes Metab Eur Heart J

Evid Rep Technol Assess (Summ)

Fam Pract Foot Ankle Int Gastroenterology Gut

Headache Health Educ Behav Health Psychol Health Technol Assess

Heart Heart Lung Int J Clin Pract Int J Nurs Stud Int J Obes (Lond)

J Adv Nurs J Allerg Clin Immunol J Am Acad Child Adolesc

Psychiatry J Am Acad Dermatol

J Am Coll Cardiol J Am Coll Surg J Am Diet Assoc

J Am Geriatr Soc J Am Soc Nephrol

J Arthroplasty J Bone Joint Surg Am J Bone Joint Surg Br J Child Psychol Psychiatry

J Clin Epidemiol J Clin Nurs J Clin Oncol

J Clin Psychopharmacol J Consult Clin Psychol J Fam Pract

J Gen Intern Med J Hand Surg [Am] J Hand Surg Eur Vol J Infect Dis

J Manipulative Physiol Ther J Neurol Neurosurg Psychiatry

J Neurosurg J Nurs Scholarsh J Orthop Trauma J Pediatr

J Pediatr Oncol Nurs

J Pediatric Orthop

J Rheumatol

J Shoulder Elbow Surg

J Trauma J Vasc Surg JAMA Kidney Int Lancet Lancet Neurol Lancet Oncology

Mayo Clin Proc Med Care Med J Aust Midwifery N Engl J Med Neurology Nurs Res

Obesity (Silver Spring) Obesity Surgery Obstet Gynecol Oncol Nurs Forum

Nutr J

Patient Educ Couns

Pediatrics Pharmacotherapy Phys Ther Prev Med

Psychosom Med Qual Health Res Radiology Res Nurs Health

Rheumatology (Oxford) Soc Sci Med Spine Spine J Stroke

Thorax West J Nurs Res

*This list is subject to modification based on the relative performance of each journal according to the set criteria.

98 EBN October 2009 Vol 12 No 4