
Evidence-Based Nursing



Join Evidence-Based Nursing Twitter Journal Chat #ebnjc

This article has been chosen by the Editor to be of specialist interest or importance

and is freely available online.



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics

www.publicationethics.org.uk





Receive regular table of contents by email. Register using this QR code.

Contents

Editorial

33 Election 2015: making sure your voice is heard

Research made simple

34 Issues of validity and reliability in qualitative research

Women's health

36 Sixteen per cent of women have depression



symptoms in the year after childbirth and are more likely to report physical symptoms, including tiredness and back pain

Midwifery

- 37 Maternal extraversion, emotional stability and conscientiousness are associated with initiation and continuation of breastfeeding
- 38 Of women seeking assistance for intimate partner violence, those who report victimisation during pregnancy are at higher risk of further victimisation
- 39 Pregnant women with schizophrenia are at higher risk of pre-eclampsia, venous thromboembolism and adverse neonatal outcomes

Adult nursing

- 41 Fasting may not be required before percutaneous coronary intervention
- 42 A decision tree incorporating biomarkers and patient characteristics estimates mortality risk for adults with septic shock
- 43 Addressing anxiety and insecure attachment in close relationships could improve quality of life for gynaecological cancer survivors
- 44 Higher red meat intake in early adulthood is associated with increased risk of breast cancer; substitution with different protein sources such as legumes and poultry may help

Child health

- 45 High consumption of peanuts or tree nuts by non-allergic mothers around the time of pregnancy reduces the risk of nut allergy in the child
- Low parent health literacy is associated with 'obesogenic' infant care behaviours
- 47 Half of parents underestimate the weight of their overweight or obese children
- 48 Cumulative adversity in early childhood is associated with increased BMI and behavioural problems

April 2015 Vol 18 No 2

- 50 Nurse home visits for infants and toddlers of low-income families improve behavioural, language and attention outcomes at age 6–9 years; paraprofessional visits improve visual attention and task switching
- 52 Individualised education reduces anxiety levels in children over 4 years undergoing elective surgery
- 53 Adoptive parents can be unprepared for the challenges in caring for children with reactive attachment disorder
- 54 Children with poor adherence to antiepileptic drugs during the first 6 months of treatment are less likely to be seizure free after 4 years

Primary healthcare

- Poor quality evidence to support the use of biofeedback for the treatment of functional constipation in adults
- A nurse-led email reminder programme on healthy lifestyle can improve cardiovascular risk factors in hypertensive adults

Pain management

- 57 Review: femoral nerve block may be the most effective option for pain relief following total knee replacement
- 58 Health professionals' pain management decisions are influenced by their role (nurse or physician) and by patient gender, age and ethnicity

Nursing issues

- 59 Infection control practice: interview with 20 nurses reveals themes of rationalising their own behaviour and justifying any deviations from policy
- Registered nurses, nurse specialists and physicians have a neutral to moderately positive view on nurse prescribing, though physicians still have patient safety concerns
- 61 High prevalence of neck, shoulder and back pain among nursing graduates warrants preventive strategies during the degree and into working life
- 62 Increasing nurse staffing levels and a higher
- proportion with bachelor's degrees could decrease patient mortality risk
- 63 Higher numbers of nurses and consultants improve survival rates for critically ill patients

Care of older people

Review: nurses predominantly have negative feelings towards the use of physical restraints in geriatric care, though some still perceive a need in clinical practice



