


Evidence-Based Nursing

July 2017 Vol 20 No 3



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc

Contents

Editorial

- 65 Is it time for a review of the way that NICE guidelines are developed, presented and used to inform policy and practice?

EBN opinion

- 67 Losing the child's voice and 'the captive mother': an inevitable legacy of family-centred care?

EBN perspective

- 70 Midwifery and women's health

Research made simple

- 74  Mixed methods research: expanding the evidence base

Women's health and midwifery

- 76 Further research supports association between antidepressant use and elevated risk of postpartum haemorrhage
- 77 Community-based exercise interventions during pregnancy are perceived as a satisfactory and motivating form of exercise engagement
- 78 Measures of depression and anxiety in women with hyperemesis gravidarum are flawed

Child health

- 80 The effectiveness of oral dextrose gel for the treatment of neonatal hypoglycaemia remains unclear
- 81  Parents and informal caregivers feel they receive insufficient communication about routine childhood vaccination

Nursing issues

- 83 Health, psychosocial and workplace characteristics may identify nurses and midwives at risk of high absenteeism
- 84 Nurses require confidence, knowledge and communication skills for referrals to doctors

- 85 Family-centred advanced care planning with adolescents living with HIV is perceived as important, helpful and meaningful
- 86 The effectiveness of protected mealtimes in hospital remains unclear
- 87 Simple variations to traditional models of care can dramatically improve emergency department performance
- 88 Intravenous paracetamol or intravenous propacetamol can provide effective postoperative analgesia for some patients
- 89  Reporting of professional misconduct is influenced by nurses' level of education and managerial experience

Care of the older person

- 91 Assessment and treatment of persistent pain in nursing home residents should be implemented systematically to prevent suffering
- 92 Cognitive-behavioural therapy for insomnia is effective, safe and highly deployable
- 93 Geriatric monitoring units could provide solution to ICU overcrowding and improve survival in older hospitalised patients
- 94 Antidepressant use associated with increased risk of hip fracture in the older population
- 95 The perspectives of patients and their families remain largely unheard in mealtime assistance research

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

COPE
Member since 2010
JN06600

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

 equator
network

 recycle
When you have finished
with this please recycle it



20 (3)

Evid Based Nurs 2017 20: 65-96

Updated information and services can be found at:
<http://ebn.bmj.com/content/20/3>

These include:

**Email alerting
service**

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>