## **Evidence-Based Nursing**

## Contents

Purpose and procedure	34	A structured and individualised safety programme reduced falls in high risk nursing home patients	52
EBN notebook	36	Seminars on advance directives for adult outpatients increased completion rates	53
Implementation forum	38	Prognosis  Grief after pregnancy loss was predicted by length of pregnancy, neuroticism, psychiatric symptoms, and absence of other children	54
Setting the results of individual studies in context	40	Causation	
Editors' response:			
Evidence-based mental health		Childhood acute lymphoblastic leukaemia was not linked to residential exposure to power line magnetic	
Editors' response:	40	fields	55
Treatment		Moderate prenatal alcohol exposure was associated with a modest increase in adolescent learning and behavioural difficulties	56
Early hospital discharge of newborn infants was not associated with feeding related hospital readmission during the first 28 days of life	41	Low birth weight was associated with non-fatal coronary artery disease and stroke in nurses	57
Early hospital discharge of newborn infants was associated with increased risk of hospital readmission during the first 28 days of life	42	Quality improvement	
Review: some psychosocial interventions can help children and families cope with chronic conditions	43	Review: telephone and computer communication improves healthcare process and patient outcomes  Reminder interventions increased women's use of	58
Telephone reminders improved compliance with a second dose of hepatitis B vaccine in high risk adults	44	mammography and Pap smear screening	59
The nicotine inhaler was an effective and safe aid to smoking cessation	45	Qualitative	
A comprehensive smoking cessation programme after surgery increased 1 year quit rates	46	Parental perception of stress during a child's health crisis was represented by 4 dimensions	60
A 2 minute surgical hand scrub was associated with higher, but not clinically significant, bacteria counts compared with a 3 minute hand scrub	47	Homeless battered women experienced patterns of violence that represented phases in their abusive	
Calcium and vitamin D reduced non-vertebral fractures		relationships	61
and bone loss in the elderly	48	Environmental issues affected women's caring	62
An empowerment information intervention improved participation in treatment decision making in men with recently diagnosed prostate cancer	49	Surgical nursing care was viewed as engaged or detached by women in hospital for hip surgery	63
Review: compression treatment improves healing of venous leg ulcers		Patients experienced physiological and psychological difficulties on long term home oxygen treatment	64
A specially designed foam mattress replacement	51	GIOSSARV Inside back co	ver