Evidence-Based Nursing

Contents

Purpose and procedure 98	Causation
Other articles noted	Pregnancy within 8 months of a previous birth was associated with an increased rate of preterm birth 122
Acknowledgements104	Higher maternal age at birth was associated with increased self sufficiency of children at age 27 to 33 years
EBN notebook	Review: parental smoking increases risk of recurrent otitis media, middle ear effusion, and tonsillectomy or
Implementation forum108	adenoidectomy in children
Treatment	Death was associated with low and high body mass index in women and older men
Offering HIV testing to pregnant women increased test uptake but not anxiety	Depression was associated with decreased survival for hospital patients who were seriously ill
Cleaning the umbilical cord with water rather than alcohol shortened the time to separation with no change in colonisation112	Quality improvement
Review: topical agents for cord care have not been shown to be effective in newborn infants in developed countries	Review: case management programmes improve patient outcomes
Review: morphine, eutectic mixture of local anaesthetics, ketorolac, and sucrose reduce postoperative and procedural pain in children114	Qualitative
A school based programme reduced dating violence in adolescents	Discovering a sense of difference was the main
Review: patient education and counselling increase	experience of growing up with cystic fibrosis 129
preventive behaviours	Deliberate HIV exposure by 6 African-American youths was rooted in growing up poor in the inner city
asthma improves health outcomes	Integrating hearing loss was a complex and dynamic process for patients
A non-prescription analgesic alleviated migraine headache pain and symptoms	Review: patients living with diabetes mellitus focus on
A 6 week psychoeducational programme reduced pain and improved quality of life in adults with chronic	learning to balance by assuming control for the management of their illness
A home based intervention reduced the frequency of hospital readmissions and out of hospital deaths after	Reimaging after an alteration in appearance or function involved 3 phases with assimilation, accommodation, and interpretation
discharge	People with drinking problems experienced suffering as a vicious, spiralling vortex that encompassed all
Diagnosis	aspects of their being
A primary healthcare team had low sensitivity for detecting postnatal depression	Glossary Inside back cover