

Evidence-Based Nursing



Contents

April 2008 Vol 11 No 2

Purpose and procedure


33

EBN notebook

- 35 Increasing research use in nursing: implications for clinical educators and managers
- 39 When should we start oral intake in children with severe acute pancreatitis?

Treatment

- 41 Nurse home visits during pregnancy and early childhood improved maternal life course 7 years later
- 42 Intranasal fentanyl and intravenous morphine did not differ for pain relief in children with closed long-bone fractures
- 43 A group intervention for parents and children achieved greater weight loss in obese children than routine care
- 44 Review: dietary interventions, with or without exercise, promote weight loss more than advice alone
- 45 Self-report, medical staff interview, and physician interview had similar effectiveness for screening for domestic violence in women
- 46 Review: delayed or immediate antibiotic prescriptions have similar clinical outcomes in respiratory infections
- 47 Review: topical permethrin was more effective than topical crotamiton or lindane for scabies
- 48 Aerobic plus resistance training was more effective than either alone for glycaemic control in type 2 diabetes
- 49 Review: evidence does not support use of static magnets for pain
- 50 Review: non-steroidal anti-inflammatory drugs and muscle relaxants are moderately effective for low back pain
- 51 Review: intensified patient care may improve adherence to lipid-lowering medication in primary or secondary prevention of CV disease

- 52 Primary prevention with pravastatin for 5 years continued to prevent coronary events in the next 10 years
- 53 Both low-dose and micro-dose 17 β -oestradiol are effective for postmenopausal hot flushes
- 54 A multicomponent cognitive-behavioural intervention for breast cancer survivors with insomnia improved perceived sleep
- 55 Review: routine NG decompression after abdominal surgery delays return of bowel function and increases pulmonary complications
- 56  Review: early feeding after major gynaecological surgery is safe except for an increase in nausea
- 57 A probiotic drink prevented diarrhoea and *Clostridium difficile*-associated diarrhoea in older patients taking antibiotics
- 58 Short-term vitamin D plus calcium did not prevent falls more than calcium alone in elderly inpatients
- 59 Review: calcium supplementation, with or without vitamin D, prevents osteoporotic fractures in people ≥ 50 years of age
- 60 A 16-week tai chi programme prevented falls in healthy older adults

Qualitative

- 61 Patients with advanced cancer used 4 self-action strategies to manage eating-related problems
- 62 Chemotherapy-induced peripheral neuropathy was described as background noise affecting daily life
- 63 Reasons for non-adherence to cardiac rehabilitation programmes included lack of motivation, domestic duties, and other health problems
- 64 Carers living with stroke survivors who were incontinent had minimal social interaction and felt socially isolated

Glossary

Inside back cover



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.



www.publicationethics.org.uk
This journal is a member of and subscribes to the principles of the Committee on Publication Ethics