


Evidence-Based Nursing



Contents

Editorial

- 31 The science and art of theoretical location

Research made simple

- 32 Selecting the sample

Women's health

- 34 85% of women with breast cancer reported changes to sexual well-being, with most wanting information on these changes
- 35 Hypertension during pregnancy is associated with increased risk of chronic and end-stage kidney disease
- 36 Hypertension during pregnancy is associated with increased risk of later cardiovascular disease, kidney disease and diabetes
- 37 Offspring of mothers who had antenatal depression and experienced maltreatment in childhood are more likely to experience child maltreatment themselves
- 38 Psychosocial and psychological interventions reduce the risk of postnatal depression compared with standard care

Adult nursing

- 40 Early ambulatory palliative care visits focus on psychosocial elements such as building rapport and coping, as well as managing symptoms and enhancing illness understanding
- 41 Five to seven years after breast cancer treatment, over a third of women (37%) report persistent pain
- 42 Small study finds that 3 years after prostate cancer treatment, men may report high quality of life and functioning
- 43 People with HIV are at increased risk of acute myocardial infarction

Child health

- 45 In China the use of analgesics and sedation following paediatric cardiac surgery is variable; average pain scores are reported to be good but over-sedation is common
- 46 Maternal alcohol-use disorder is associated with increased risk of sudden infant death syndrome and infant death from other causes

April 2014 Vol 17 No 2

Primary healthcare

- 48 Daily calcium intake in excess of 1400 mg is associated with increased all-cause and cardiovascular disease mortality in women
- 49 Prime time youth development intervention improves contraceptive use and sexual awareness among sexually active adolescent girls

Pain management


- 51 Limited evidence that single-dose oral ibuprofen plus codeine is more effective for postoperative pain than either drug alone

Nursing issues

- 53 In intensive care and bone marrow transplantation settings, daily bathing with chlorhexidine wash cloths reduces the risk of hospital-acquired infection
- 54 Nine per cent of nurses across Europe report intent to leave their profession, with burnout among the associated personal and professional factors
- 55 Modelling suggests authentic leadership from managers influences structural empowerment, job satisfaction and self-rated performance among nurses
- 57 Use of multifaceted strategies and processes at multiple levels, and passionate frontline leaders are associated with uptake of best practice guidelines on breastfeeding

Care of the older person

- 59 Older adults reporting social isolation or loneliness show poorer cognitive function 4 years later
- 60 Community-dwelling older adults with hearing loss experience greater decline in cognitive function over time than those with normal hearing
- 61 Prescriber preference may influence prolonged antibiotic usage more than patient characteristics in long-term care facilities

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it



Receive regular table of contents by email.
Register using this QR code.