## **Evidence-Based Nursing**

## Contents

Purpose and procedure	66	Bladder training plus pelvic muscle exercises reduced urinary incontinence in women immediately after treatment.	85
EBN notebook		A nurse led programme improved physical function in older adults with chronic disease	86
If you could just provide me with a sample: examining sampling in qualitative and quantitative research papers	68	Quality improvement	
Implementation forum		Review: nursing care driven by guidelines improves some process measures and patient outcomes	87
Continuing professional development in Canada and the UK: how evidence-based resources can help	71	Causation	
Treatment		Frequent nut consumption reduced the risk of coronary heart disease in women	88
Antenatal oral dexamethasone for neonatal respiratory		Dietary fibre intake was not associated with a lower risk of colorectal cancer in women	89
distress syndrome increased neonatal morbidity	73	Multiple attempts to stop smoking, even with relapses, reduced loss of pulmonary function	90
Review: non-nutritive sucking decreases length of hospital stay in premature infants	74	reduced loss of pullionary function	30
Review: support interventions reduce cessation of breast feeding before 2 months	75	Economics	
Soy milk decreased chronic constipation in young children	76	Early opportunistic screening was cost effective in young adults with type 2 diabetes	91
Review: antibiotics are not effective for upper respiratory tract infection in children	77	Qualitative	
Review: antibiotics did not lead to general improvement in upper respiratory tract infections	78		
Review: controlling house dust mites is ineffective for asthmatic patients who are sensitive to mites	79	During cervical screening, women desired good communication with their doctor and to be treated with sensitivity	92
Review: self help interventions alone minimally increase smoking cessation rates	80	Women's decisions to seek evaluation of self discovered breast symptoms occurred in a complex social context .	93
Review: minor analgesics are effective for primary dysmenorrhoea	81	Patient distress after radiotherapy to the head and neck was often hidden	94
Review: St John's wort is more effective than placebo for mild to moderate depressive disorders	82	Fatigue in HIV infection was perceived as a barometer of disease progression	95
Individualised computer generated nutrition information plus iterative feedback reduced dietary fat and increased fruit and vegetable intake	83	Patients focused on integrating diabetes into their daily lives; practitioners focused on glucose control	96
Pressure bandages after coronary angiography reduced bleeding, but increased discomfort	84	Glossary	over