Evidence-Based Nursing

Contents

Purpose and procedure	2	Glargine dose titration by patients and by physicians were equally effective for preventing severe	20
EBN notebook		hypoglycaemia	20
Helping practitioners understand the contribution of qualitative research to evidence-based practice	4	Web based care management improved glucose control in patients with poorly controlled diabetes	21
Reflections on "Helping practitioners understand the contribution of qualitative research to evidence-based practice"	7	An educational booklet did not improve adherence or symptoms in patients prescribed thyroxine for primary hypothyroidism	22
Treatment		Review: multidisciplinary interventions reduce hospital admission and all cause mortality in heart failure	23
The WHO technique for intramuscular thigh vaccination in infants and toddlers had fewer adverse reactions than 2 other techniques	9	Review: multivitamins and mineral supplements do not reduce infections in elderly people	24
Liposomal lidocaine improved intravenous cannulation success rates in children	10	Multivitamin and multimineral supplements did not reduce reported infection days or related use of healthcare services in elderly people	25
Review: children permitted clear fluids ≤ 120 minutes before surgery have similar gastric volumes and pH		Clinical prediction guide	
values as those on standard fasts	11	A simple risk score predicted 7 day stroke risk after transient ischaemic attack	26
Topical chloramphenicol was not effective for acute infective conjunctivitis in children	12	Qualitative	
Nurse home visits did not differ from standard care for prevention of recurrent child abuse	13	Mothers' decisions about MMR vaccination were framed by their children's vulnerabilities and wider social trends	27
Multisystemic therapy improved adherence to blood glucose testing in adolescents with type 1 diabetes	14	UK and US adolescents perceived internet health information to be salient but of questionable credibility	28
A weight maintenance diet reduced bulimic symptoms in adolescent girls	15	Young women described the benefits of having advance supplies of emergency contraception but emphasised its	
After 3 months, low dose oral contraceptives reduced pain in adolescent girls with moderate to severe	17	use as a "last resort" rather than an alternative form of contraception	29
dysmenorrhoea Trimethoprim reduced dysuria in women with	16	Marginalised HIV positive drug users felt that an HIV diagnosis created benefits as well as losses	30
symptoms of urinary tract infection but negative urine dipstick test results	17	Perceived barriers and benefits were factors in decision making about colorectal screening	31
Review: wearing graduated compression stockings during air travel reduces the risk of deep venous thromboembolism	18	UK palliative care professionals identified service infrastructure, patient and carer attitudes and characteristics, and practice culture as influencing place of death of patients with cancer	32
A home based, physical activity intervention increased physical activity, fitness, and vigour and reduced fatigue in sedentary women with early stage breast cancer	19	Glossary Inside back co	

Search all issues of *Evidence-Based Nursing* published since 1998 and over 300 medical journals through Highwire press at www.evidencebasednursing.com. EBN 13845.