

Evidence-Based Nursing

Contents

Purpose and procedure.....	66	Children who were sexually abused were more disturbed than their peers after 5 years	84
EBN notebook.....	68	Causation	
Implementation forum.....	71	Review: previous caesarean delivery or abortion increases the risk of placenta praevia	85
Treatment		Review: parental tobacco smoke increases the risk of asthma and respiratory symptoms in school age children	86
Early discharge after nurse specialist education and planning plus follow up visits and telephone calls was cost effective for high risk pregnant women and their newborn infants	73	Review: use of hormone replacement therapy in the past 5 years is associated with an increased risk of breast cancer	87
Review: sucrose reduces crying during painful procedures in newborn infants	74	Review: smoking is associated with decreased bone mineral density and increased risk of hip fracture in postmenopausal women	88
Review: non-parental daycare has short term and long term benefits for preschool children	75	Emotional support and mastery were associated with decreased mortality in older people	89
Nurse home visits to low income, first time mothers reduced pregnancy induced hypertension and childhood injuries	76	Clinical prediction guide	
Nurse home visits reduced child abuse and neglect over a 15 year period	77	A risk assessment score predicted which elderly patients would fall during a hospital stay	90
Return of school forms and nurse home visits increased adherence with follow up reading of tuberculosis tests in children	78	Qualitative	
Stress point interventions for parents of children in hospital with chronic conditions reduced stress and improved child and family functioning	79	Nurses responded on an ad hoc basis to the unpredictable needs of parents of children in hospital ...	91
A social support intervention alone and with reminders improved appointment keeping among patients with chronic illnesses	80	Mothers of ill children experienced uncertainty before asthma was diagnosed in their children	92
Fewer patients dislodged peripheral intravenous catheters with transparent dressings than with gauze dressings	81	Families with children diagnosed with cancer used various strategies to create a <i>new normal</i> routine	93
As needed in line suction catheter changes were as safe as and less expensive than daily scheduled catheter changes during mechanical ventilation	82	The experience of giving up smoking was described as encompassing 7 major categories	94
Prognosis		Patients' experiences of healing a venous ulcer were described as "forever healing" which affected their activities and treatment	95
Unintentional pregnancy was associated with reduced intentions to breast feed	83	Older adults worked through 3 phases as they made the initial transition to permanent residence in a nursing home	96
		Glossary	Inside back cover