Evidence-Based Nursing

Contents

Purpose and procedure	66	Review: acupuncture and mind-body therapies may be effective for cancer related pain	84
EBN notebook		A proactive communication strategy reduced post-	
The synthesis of qualitative research and evidence-based nursing	68	traumatic stress disorder symptoms in relatives of patients dying in the ICU	85
Teaching tip: using the "Who wants to be a millionaire?" game to teach searching skills	72	Overnight splinting of the wrist in a neutral or extended position did not prevent contracture after stroke	86
Treatment	, 2	4 layer bandages were better than 1 layer bandages, and pentoxifylline may be better than placebo for venous leg ulcers	87
Review: antihistamines, decongestants, or both do not provide benefit in children with otitis media with effusion	73	Daily folic acid supplementation for 3 years reduced age related hearing loss	88
Delivery of a β_2 agonist by metered dose inhaler with a bottle spacer was equivalent to delivery by conventional spacer in young children with acute lower airway obstruction	74	An exercise programme led to a slower decline in activities of daily living in nursing home patients with Alzheimer's disease	89
Review: distraction, hypnosis, and combined cognitive- behavioural interventions reduce needle related pain and distress in children and adolescents	75	Causation Weak infant sucking in the first 8 weeks, reliance on breastfeeding for ≥9 months, small feeds, and difficulties	
Review: over-the-counter medications are effective for gastro-oesophageal reflux disease	76	in weaning were associated with failure to thrive	90
The LiftTrainer TM programme was not more effective than a video for reducing back injury in jobs with repetitive lifting	77	Quality improvement Including a weaning predictor in the daily assessment of weaning readiness increased mechanical ventilation weaning time	91
Review: lifestyle or pharmacological interventions prevent or delay type 2 diabetes in people with impaired glucose tolerance	78	Qualitative Clinically obese children identified facilitators and	
Review: oral anticoagulants plus aspirin reduce arterial thromboembolism more than oral anticoagulants alone		barriers to initiating and maintaining the behaviours required for weight loss	92
but only in patients with mechanical heart valves Review: adequately randomised trials showed that mammography screening did not significantly reduce	79	Children were marginalised by parents and healthcare professionals during consultations for dietary management of cystic fibrosis	93
breast cancer, cancer, or all cause mortality but increased breast surgeries	80	Lesbian women and gay men found that nurses often assumed they were heterosexual, which led to feelings	0.4
Oral nutritional supplements during acute illness and recovery reduced non-elective hospital readmissions in older patients	81	of discomfort and insecurity A diagnosis of HIV infection evoked strong and long lasting emotional reactions in women	94 95
Review: timing or administration route of nutritional support does not affect death or disability in head injury	82	Patients with recently diagnosed hypertension described risk in terms of acceptance and denial narratives, which served as personal frameworks of coping	96
Review: patient controlled opioid analgesia reduces postoperative pain more than conventional as-needed opioid analgesia	83	Glossary Inside back to	

Search all issues of *Evidence-Based Nursing* published since 1998 and over 300 medical journals through Highwire press at www.evidencebasednursing.com. EBN 98221.