

Evidence-Based Nursing

Contents

Purpose and procedure	1	Review: several interventions reduce fear of falling in older people living in the community	21
EBN notebook		In-home palliative care increased patient satisfaction and reduced use and costs of medical services.....	22
New Year—New Look!	3		
The use of qualitative evidence in clinical care.....	4	Assessment (screening or diagnosis)	
Putting evidence into context: some advice for guideline writers	6	Review: some screening tests for dementia are accurate and practical for use in primary care	23
Treatment		Causation	
Review: sexual abstinence only programmes do not affect STIs or HIV risk behaviours in high-income countries ...	9	Children who consumed sugar-sweetened beverages between meals ≥ 4 –6 times/week at 2.5–4.5 years of age were more likely to be overweight at 4.5 years of age .	24
Starting oral contraceptives immediately or during next menses did not differ for continued contraceptive use or pregnancy at 6 months.....	10	Quality improvement	
Review: vaccines against human papillomavirus prevent cervical cancer precursors in young women	11	Cleaning of the perineal/genital area before urine collection from toilet-trained children prevented sample contamination.....	25
Sequential counselling that targeted individual health behaviours did not differ from simultaneous counselling that targeted multiple behaviours.....	12	Patient education after inguinal hernia surgery did not differ from routine information for pain at rest at 7 days	26
A tailored internet intervention did not increase physical activity more than a tailored print intervention or publicly available web sites	13	Qualitative	
Review: dietary restriction, with or without aerobic exercise, promotes weight loss in postpartum women .	14	Smoking cessation was difficult for adolescents because of daily life stressors and the need for major lifestyle changes	27
Review: regular brisk walking improves cardiovascular risk factors in healthy sedentary adults.....	15	The experience of injecting drugs in public spaces was characterised by urgency, a need for privacy, hygienic concerns, and a sense of shame	28
An intensive lifestyle intervention reduced weight and cardiovascular disease risk factors in overweight and obese people with type 2 diabetes	16	Staring, tone of voice, anxiety, mumbling, and pacing in the ED were cues for violence toward nurses	29
Review: moderate weight loss reduces functional disability but does not reduce pain in obese patients with knee osteoarthritis.....	17	Inpatients with cerebral palsy and complex communication needs identified barriers to communicating with nurses	30
Review: intensive behavioural counselling interventions are effective for smoking cessation in patients admitted to hospital.....	18	Women's experiences of myocardial infarction were described in terms of gradual onset, not having chest pain, and responding to symptoms	31
Review: effectiveness of intensive case management for severe mental illness depends on baseline hospital use and organisation.....	19	Middle-aged people with chronic heart failure experienced the frustrations of living with an unpredictable and failing body.....	32
Review: patients in nursing-led units are better prepared for discharge than those receiving usual care.....	20	Glossary	Inside back cover

Search all issues of *Evidence-Based Nursing* published since 1998 and over 300 medical journals through Highwire press at www.evidencebasednursing.com. EBN 59763.