


# Evidence-Based Nursing



## Contents

October 2008 Vol 11 No 4

### Purpose and procedure

97

### EBN notebook

- 99 Critical appraisal of cost-effectiveness and cost-utility studies in health care
- 103 Evidence-based practice targets the individual patient. Part 1: how clinicians can use study results to determine optimal individual care

### Treatment

- 105 Review: home-made and commercial spacers did not differ for delivery of inhaled  $\beta_2$  agonists in children with asthma or wheezing
- 106 A needle-free powder lignocaine delivery system reduced the pain of venipuncture in children
- 107 Topical aqueous 2% lignocaine eardrops reduced ear pain in children with acute otitis media
- 108 Review: contraceptive patch and vaginal ring are as effective as oral contraceptives
- 109 Review: evidence on the effectiveness of interventions to improve patient adherence to prescribed medications is limited
- 110 Review: screening or case-finding questionnaires used alone are not effective for management of depression
- 111 Review: preprocedure information, breast cushions, and patient-controlled breast compression reduce mammography pain
- 112 Review: reflexology improved urinary symptoms in multiple sclerosis but was not effective for other outcomes in various conditions
- 113 An automated external defibrillator in the home did not reduce all-cause mortality in patients at risk of cardiac arrest
- 114 Review: some interventions may reduce catheter-related bloodstream infections and colonisation in the ICU
- 115 Review: evidence of the effectiveness of hydrocolloids for healing pressure ulcers is limited
- 116 Negative pressure wound therapy promoted healing of diabetic foot ulcers more than advanced moist wound therapy
- 117 Review: low-molecular-weight heparin prevented recurrent VTE more than oral anticoagulants in patients with cancer

118 Review: walking aids, chest-wall vibration, and neuroelectrical muscle stimulation relieve breathlessness in COPD



- 119 Automatic stop orders reduced duration of indwelling urinary catheterisation in hospital
- 120 A multifactorial intervention did not prevent falls or fractures in elderly patients during short hospital stays

### Assessment (screening or diagnosis)

- 121 Review: simple questions and clinical tests are moderately useful for diagnosing urinary incontinence

### Prognosis

- 122 Chorioamnionitis, gestational age, male sex, birth weight, and illness severity predicted positive autism screening scores in very-low-birth-weight preterm infants

### Causation

- 123 Enalapril and captopril increased mortality compared with ramipril in elderly patients with heart failure

### Quality improvement

- 124 Protocol-directed sedation did not reduce duration of mechanical ventilation or hospital stay in ICU patients

### Qualitative

- 125 In teenagers with diabetes, transition toward autonomy in self-management involved growth in self-reliance and support from others
- 126 4 themes described the experiences of patients before, during, and immediately after awake craniotomy
- 127 Many patients with coronary heart disease believed that stress and lifestyle factors caused their illness
- 128 Patients with terminal illness and their healthcare providers preferred a dosed and gradual process of truth-telling

### Glossary

Inside back cover



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics  
[www.publicationethics.org.uk](http://www.publicationethics.org.uk)