


Evidence-Based Nursing



Contents

Purpose and procedure

1

EBN notebook

- 3 To our future—welcome students!
- 4 To lower or not to lower? Making sense of the latest research on intensive glycaemic control and cardiovascular outcomes
- 7 Applying evidence: what's the next action?

Treatment

- 9 Mothers were less likely to be depressed after a structured behavioural intervention for infant sleep problems
- 10 Review: varenicline, bupropion, and nicotine replacement therapies are effective for smoking cessation at 6 or 12 months
- 11 Review: exercise plus diet prevents type 2 diabetes
- 12 Intensive glucose control did not prevent important complications in type 2 diabetes
- 13 Intensive glucose control increased mortality and did not prevent cardiovascular events in type 2 diabetes
- 14 Benefits of early intensive glucose control in preventing diabetes-related complications were sustained for up to 10 years
- 15 Review: culturally appropriate health education improves glycaemic control in members of ethnic minority groups with diabetes
- 16 Review: short-term medical use of cannabis increases risk of non-serious adverse effects
- 17 A nurse-delivered intervention was effective for depression in patients with cancer
- 18 Review: multicomponent educational interventions improve some outcomes in predialysis and dialysis care
- 19 Clinically indicated and routine replacement of peripheral intravenous catheters did not differ for catheter failure
- 20 Review: chewing gum after abdominal surgery reduces time to return of bowel function but not length of hospital stay

21



Normal food at will and nil-by-mouth enteral feeding after major upper GI surgery did not differ for mortality or morbidity

January 2009 Vol 12 No 1

- 22 Initial refeeding with a soft diet was better than a clear liquid diet for length of hospital stay in mild acute pancreatitis
- 23 Review: psychological interventions prevent depression after stroke
- 24 Review: providing information improves subjective outcomes but may not improve clinical outcomes in patients with stroke or their carers
- 25 General practitioners' advice to use topical rather than oral ibuprofen resulted in equivalent effects on chronic knee pain

Prognosis

- 26 Excess mortality after HIV seroconversion has greatly decreased in the past 10 years

Quality improvement

- 27 Review: quality improvement interventions reduce unnecessary prescription of antibiotics for outpatients

Qualitative

- 28 Men were surprised by the severity of symptoms they experienced after laparoscopic radical prostatectomy
- 29 Patients felt greater personal control and emotional comfort in hospital when they felt secure, informed, and valued
- 30 Prioritising mealtime care, patient choice, and nutritional assessment were important for older inpatients' mealtime experiences
- 31 Peer relationships were important to the experience of thriving in some, but not all, nursing home residents
- 32 Many older people felt that electronic care surveillance increased their safety and enabled them to live alone in their own homes

Glossary

Inside back cover



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk