


Evidence-Based Nursing



Contents

Purpose and procedure

Aetiology

- 101** Use of oral contraceptives is not associated with increased risk of death in the UK; a net benefit in all-cause mortality was seen in ever users versus never users – RR 0.88, 95% CI 0.82 to 0.93

Healthcare economics

- 102** Increasing nurse staffing levels in cardiac surgery centres appears to be a cost effective patient safety intervention

Prevalence

- 103** Nineteen per cent of paediatric inpatient medication orders were associated with administration errors and 13.2% had prescribing errors in five London hospitals
- 104** National survey shows the majority of nurses use very little research in the first 2 years after their graduation, highlighting a gap between research and clinical practice
- 105** US survey finds higher availability of palliative care programs, palliative physicians and consultation teams and palliative outpatient services in National Cancer Institute centres compared to non-NCI centres
- 106** In surveyed Australian medical-surgical units about a third of nurses on perceive emotional abuse in recent shifts and a fifth report actual violence; perception of violence in the workplace is associated with unstable or negative working conditions and adverse patient outcomes
- 107** Severe adverse reactions are rare with infusions of infliximab in a community setting
- 108** Survey finds only 15% of emergency departments have formal alcohol screening and intervention policies for trauma patients

Prognosis

- 110** Activity levels inversely associated with weight gain in women only when BMI is less than 25
- 111** Nurses rate readiness for discharge higher than patients do, and nurses scores predict readmission or ED utilisation after discharge better than patient self-assessment

Qualitative

- 113** Survey of people with type 2 diabetes shows the majority prefer diabetes education to be given during regular diabetes check-ups; other education preferences and opinions on self-care vary depending on patient characteristics

October 2010 Vol 13 No 4

- 114** Limited literature available regarding the role of nurses, midwives and health visitors in development and implementation of protocol-based care
- 115** Different attitudes towards mental health revealed in a survey of nurses across five European countries; more positive attitudes found in Portugal, in women and in those in senior roles
- 116** Small survey finds that new mothers use self-care approaches to manage fatigue more often than other strategies such as seeking help from others

Quality improvement

- 118** Computer-assisted module for nursing students provides similar improvements in handwashing knowledge and may improve handwashing practice compared with face-to-face teaching

Therapeutics

- 120** Antimicrobial silver dressings beneath compression for venous ulceration are not cost-effective compared with standard dressings
- 121** A web-based tailored educational intervention increases positive perceptions of mammography and intention to obtain mammography compared to standardized mammography brochures among Taiwanese women
- 122** A nurse-led disease management programme for chronic kidney disease improves outcomes such as adherence to diet and continuous ambulatory peritoneal dialysis
- 123** Administration of annual oral high-dose vitamin D to community dwelling older women in autumn and winter months increases risk of falls and fractures
- 124** Use of a project nurse to encourage evidence-based falls injury prevention strategies did not reduce falls or fall injuries over 17 months in residential aged care facilities in Australia
- 125** Nurse-delivered, home-based pragmatic rehabilitation has a short-term effect on improving fatigue in people with chronic fatigue syndrome compared with usual GP care, but effects were not sustained at 1 year
- 126** 57% of RCTs of cardiovascular nursing interventions show that they improve at least one outcome for secondary prevention patients; optimum intervention strategy unclear

Treatment

- 128** Care management increases the use of primary and medical care services by people with severe mental illness in community mental health settings

Republished from DTB

- 130** Update on treatments for head lice



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

equator network

recycle

When you have finished with this please recycle it