


# Evidence-Based Nursing



## Contents

July 2011 Vol 14 No 3

### Editorial

- 63 Using research findings in nursing practice

### Research made simple

- 64 What is a systematic review?

### Adult nursing

- 65 Psychosocial and information needs of people affected by colorectal cancer are poorly met during follow-up
- 66 Listening to music may relax mechanically ventilated patients, but there are limitations to the quality of the available evidence
- 68 A 12-month intensive supervised exercise intervention and counselling reduces HbA<sub>1c</sub>, blood pressure and other modifiable cardiovascular risk factors in people with type 2 diabetes
- 69 A characterisation of factors determining postoperative ileus after laparoscopic colectomy enables the generation of a novel predictive score
- 71 Taking aspirin daily for at least 4 years reduces long-term risk of cancer death
- 72 In people with poorly controlled type 1 diabetes, cognitive behaviour therapy combined with motivational enhancement therapy reduces HbA<sub>1c</sub> after 12 months

### Care of the older person

- 74 Pooled results of two small trials with few events show a non-significant trend towards increased risk of intracerebral haemorrhage in people with Alzheimer's disease
- 75 Avoid benzodiazepines and opioids in people at risk of delirium
- 76 The Otago strength and balance exercise programme lowers the risk of death and falls in the older people at 12 months
- 78 People with Alzheimer's disease are at increased risk of hip fracture and of mortality after hip fracture
- 79 Skin protection wheelchair cushions for older nursing home residents reduce 6-month incidence of ischial tuberosity pressure ulcers compared with segmented foam cushions

### Child health

- 81 Efficacy of implementation intentions intervention on prevention of smoking among adolescents

### Mental health

- 83 Assessment following self-harm: nurses provide comparable risk assessment to psychiatrists but are less likely to admit for in-hospital treatment

### Midwifery

- 85 For pregnant women near full term with intrauterine growth restriction, induction of labour does not affect risk of adverse neonatal outcome or caesarean section compared with expectant management

### Nursing Issues

- 87 An evidence-based communication skills training programme for oncology nurses improves patient-centred communication, enhancing empathy, reassurance and discussion of psychosocial needs
- 88 The organisational culture of nursing staff providing long-term dementia care is related to quality of care

### Pain management

- 90 Fentanyl Pectin Nasal Spray reduces breakthrough cancer pain intensity compared with placebo in people taking at least 60 mg daily oral morphine or equivalent

### Women's health

- 92 Incorporating mental healthcare into routine postpartum nursing is a promising approach for reducing depression in new mothers

### Resources page

- 94 EBN Resources page


### Electronic page

- e1 Purpose and procedure

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

**C O P E** COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics  
[www.publicationethics.org.uk](http://www.publicationethics.org.uk)

 equator network

 recycle  
When you have finished with this please recycle it