


# Evidence-Based Nursing



## Contents

January 2012 Vol 15 No 1

### Editorial

- 1 What is quality in qualitative health research?

### Research made simple

- 2 Qualitative data analysis



### Health promotion

- 3 In smokers not willing to quit, counselling on smoking reduction plus free nicotine replacement therapy, compared with one-off cessation advice, increases the proportion achieving abstinence or reduction in smoking rate at 6 months

### Nursing issues

- 5 Three-year follow-up after introduction of Canadian best practice guidelines for asthma and foot care in diabetes suggests that monitoring of nursing-care indicators using an electronic documentation system promotes sustained implementation
- 6 Review of research findings suggesting nurses overreport their use of research
- 7 Use of real-time continuous glucose monitoring versus traditional self-monitoring of blood glucose levels improves glycaemic control in patients with type 1 diabetes
- 9 Best interest standards do not correlate with the reality of physicians' decision making in life and death choices

### Midwifery

- 10 Pregnant women who experienced late stillbirth appear less likely to have slept on their left
- 11 2009/H1N1 infection in pregnancy association with adverse perinatal outcomes

### Child health

- 13 Installation of safety devices reduces the risk of home injury in children
- 14 Provision of essential newborn care training to midwives in Zambia is a low-cost intervention that reduces neonatal mortality
- 15 Nurse:patient ratios influence the achievement of oxygen saturation targets in premature infants

### Primary healthcare

- 17 Diabetes management can be safely transferred to practice nurses
- 18 Nurse case management with a therapeutic algorithm for people living with diabetes, hypertension and raised LDL cholesterol: after 1 year 22% of those receiving the intervention have all three parameters under control versus 10% of those receiving usual care

### Care of the older person

- 20 Social aspect of activity stimuli is related to positive affect in persons with Alzheimer's disease
- 21 End-of-life care in two Norwegian nursing homes: family perspectives
- 22 In previously continent adults, aged 70 or older, use of urinary catheters or diapers while in hospital increases the risk of new urinary incontinence
- 23 Reviewing the evidence base for nurse staffing and quality of care in nursing homes
- 25 Review: comprehensive geriatric assessment increases a patient's likelihood of being alive and in their own home at up to 12 months
- 26 Improving the ability to predict falls among older adults following inpatient rehabilitation

### Adult nursing

- 28 Relationships between patients, informal caregivers and health professionals in care homes
- 29 Complementary and alternative medicine is popular among chronic renal failure patients – renal teams must increase their competence to advise patients with respect to efficacy and safety
- 30 Experiences of 12 patients in the first 2 years following spinal cord injury: setting meaningful rehabilitation goals
- 32 Racial Disparities in the Incidence of Diabetes in Canada

### Electronic page

- e1 Purpose and procedure



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics  
[www.publicationethics.org.uk](http://www.publicationethics.org.uk)

equator network

recycle

When you have finished with this please recycle it