


Evidence-Based Nursing



Contents

Editorial

- 99 Can we help patients have a better experience?
Implementing NICE guidance on patient experience

Research made simple

- 101 What is a qualitative synthesis?



Midwifery

- 102 Maternal perception of altered fetal activity and the risk of late stillbirth
- 103 Perineal warm compress reduces risk of third- and fourth-degree tears and should be part of second stage care
- 104 Major depression in pregnancy and post partum associated with anxiety disorders and substance use
- 105 Cochrane review: relaxation and yoga may decrease pain during labour and increase satisfaction with pain relief, but better quality evidence is needed
- 106 Cochrane review finds that use of cardiotocograph on admission to the labour ward, rather than intermittent auscultation of the fetal heart rate, may increase risk of caesarean in low-risk women

Adult nursing

- 108 Systematic review and meta-analysis of the correlates of cancer-related fatigue
- 109 Adding targeted multiple interventions to standard fall prevention interventions reduces falls in an acute care setting
- 110 In Israel, most families of cadaver organ donors and transplant recipients who had made contact wanted transplant coordinators to facilitate this contact

Child health

- 112 Interviews of 21 adolescents reveal a desire for structured education programme specifically tailored to their needs
- 113 TV viewing, but not total sedentary behaviour, is associated with adverse cardiometabolic biomarkers in adolescents
- 114 No evidence that introduction of allergenic foods prior to 6 months of age increases reports of wheeze or eczema in young childhood
- 115 Review finds that bed sharing increases risk of sudden infant death syndrome



October 2012 Vol 15 No 4

- 116 Youth physical activity and sedentary time and associations with cardiometabolic health
- 117 There is no evidence to suggest that a computerised individually-tailored intervention prevents weight gain in adolescents

Health promotion

- 119 Nurse-led central venous catheter insertion: review of 760 procedures performed across three hospitals reveals a low rate of complications
- 120 In people aged over 45, increased time spent sitting daily is associated with increased risk of all-cause mortality independent of physical activity level
- 121 Higher red meat consumption is associated with increased risk of all-cause, cardiovascular, and cancer mortality

Care of older people

- 123 Interviews with Australian nursing home managers reveal that various factors are associated with the decision to transfer a resident to hospital, including two factors not identified in previous research: advance care planning and support from local health services

Mental health

- 125 The reciprocal relationship between physical activity and depression in older European adults



Pain management

- 126 Administering enteral tube medications – room for improvement?
- 127 Review of variable quality trials finds that warming local anaesthetic to body temperature reduces self-reported pain of infiltration by 11 mm on a 100 mm rating scale

Miscellaneous

- 128 Correction

Electronic page

- e4 Purpose and procedure

This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.



C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

equator network

recycle

When you have finished with this please recycle it