


Evidence-Based Nursing

July 2013 Vol 16 No 3



Contents

Editorial

- 69 Patient and public involvement in research: future challenges

Research made simple

- 70 Blinding: an essential component in decreasing risk of bias in experimental designs



Resources

- 72 EBN resources page

Midwifery

- 74 Use of antidepressants in the second trimester is associated with reduced pregnancy duration, and third trimester antidepressant use with infant convulsions
- 75 37% of child survivors of intrauterine or neonatal insults experience at least one long-term sequela, the most common being neurodevelopmental delay
- 76 Women who drink heavily during pregnancy have increased stillbirth risk

Mental health

- 78 Self-harm within inpatient psychiatric services: most episodes are among women, involve breaking the skin and take place in private
- 79 In individuals with schizophrenia, duration and severity of mental illness, health status, mutuality, employment and household income influence quality of life

Adult nursing

- 81 Among patients admitted to ICU, delirium is most common in those with neurological diagnoses, and is associated with adverse health outcomes
- 82 Postangiography patients who complete cardiac rehabilitation have reduced risk of mortality or hospitalisation compared to non-completers
- 83 Postmenopausal women with medically treated diabetes have increased risk of lung cancer
- 84 Early ambulation after percutaneous coronary intervention does not increase bleeding risk compared with late ambulation
- 86 Predicting outcome at the bedside: the surgical intensive care unit: optimal mobility score

Child health

- 87 Adolescents who take part in team sports, or who actively commute to school, are less likely to be obese

Primary healthcare


- 89 Quality improvement strategies for diabetes management decrease HbA1c, cholesterol and blood pressure, and increase screening for disease complications

Nursing issues

- 91 The majority of nurses first experience death in or before their first year of practice, and the experience can provoke feelings of helplessness, guilt and ongoing distress
- 92 Aspirin given for up to 2 years after initial anticoagulant treatment reduces the risk of venous thromboembolism recurrence without increasing risk of major bleeding

Care of older people

- 94 Compression improves healing of venous leg ulcers compared with no compression, with differences between different compression systems
- 95 Older people who report loneliness have increased risk of mortality and functional decline

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

 equator network

 recycle

When you have finished with this please recycle it



Receive regular table of contents by email.
Register using this QR code.