


Evidence-Based Nursing



Contents

July 2014 Vol 17 No 3

Editorial

- 63 Disclosure of domestic violence and abuse: how prepared are you?

Research made simple

- 65  Service evaluation, audit and research: what is the difference?

Resources

- 67 EBN resources page

Women's health

- 71 Drospirenone-containing oral contraceptives may increase the risk of venous thromboembolism
- 72 In women, increased dietary antioxidants are associated with reduced risk of developing heart failure
- 73 Available evidence suggests women with false-positive mammograms less likely to return for screening

Midwifery

- 74 Second trimester hyperemesis gravidarum is associated with increased risk of preterm pre-eclampsia, placental abruption and small for gestational age birth
- 75 Gestational diabetes: higher animal protein intake during pregnancy is associated with increased risk, and higher vegetable protein intake with decreased risk
- 76 Late pregnancy use of selective serotonin reuptake inhibitors and serotonin and norepinephrine reuptake inhibitors is associated with increased risk of postpartum haemorrhage
- 77 Antenatal perineal massage decreases risk of perineal trauma during birth
- 78 In Western Australia, 70% of mothers of babies with fetal alcohol syndrome did not have an alcohol-related diagnosis recorded during pregnancy
- 79 Adolescent girls who experience abuse or neglect are at an increased risk of teen pregnancy

Mental health

- 80  People with schizophrenia are more likely to have a mother who smoked during pregnancy than people without the condition

Adult nursing

- 81  Home palliative care services increase the chance of adults with advanced illness dying at home and reduce symptom burden without impact on caregiver grief

Child health

- 82 Mandatory gloving in acute care paediatric units associated with decreased risk of hospital-acquired infections
- 83 Parents of British Bangladeshi children undergoing cancer treatment report managing competing knowledge, vigilance, advocacy, balancing work and treatment burden impacts on their lives

- 84 Prenatal exposure to sodium valproate is associated with increased risk of childhood autism and autistic spectrum disorder
- 85 Excess weight gain in the first 18 months of life is associated with later childhood overweight, obesity and greater arterial wall thickness
- 86 Children of mothers with iodine deficiency during pregnancy are more likely to have lower verbal IQ and reading scores at 8–9 years of age
- 87 Alcohol consumption in early and late pregnancy is associated with poor child reading and writing ability at 8–9 years of age
- 88 Children's experience of postoperative pain relief: children, parents and nurses use various pharmacological and non-pharmacological approaches, particularly distraction

Care of people with learning disabilities

- 89 In adults with intellectual disability, discontinuation of antipsychotics is associated with reduction in weight, BMI, waist circumference and blood pressure

Primary health care

- 90 Calcium supplements associated with increased risk of cardiovascular death in men but not women
- 91 One year following injury, pain and physical factors affect the likelihood of residual disability, but psychiatric symptoms may have a greater influence
- 92 Modest salt intake reduction for 4 weeks or longer decreases blood pressure

Nursing issues

- 94 End-of-life care in the emergency department: nurses who invest in the nurse–patient relationship are better able to manage the emotional aspects of caring for dying people and their relatives
- 95 Terminally ill patients who are supported by religious communities are more likely to receive aggressive end-of-life care rather than hospice care; spiritual support from medical teams may reverse this
- 96 The majority of hospitalised elderly people at high risk of dying have thought about end-of-life care, though documentation of preferences in medical records may be lacking
- 97 Women's experience of abuse in a healthcare setting reveals preceding loss of power struggles, with sensitivity and dependency making them vulnerable to staff domination

Care of the older person

- 98 A transitional care model for low-income older adults does not reduce readmission rates or emergency department visits during care transitions

PostScript

- 66 Correction

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it



Receive regular table of contents by email.
Register using this QR code.