


# Evidence-Based Nursing




## Contents

July 2014 Vol 17 No 3

### Editorial

- 63 Disclosure of domestic violence and abuse: how prepared are you?

### Research made simple

- 65  Service evaluation, audit and research: what is the difference?

### Resources

- 67 EBN resources page


### Women's health

- 71 Drospirenone-containing oral contraceptives may increase the risk of venous thromboembolism
- 72 In women, increased dietary antioxidants are associated with reduced risk of developing heart failure
- 73 Available evidence suggests women with false-positive mammograms less likely to return for screening


### Midwifery

- 74 Second trimester hyperemesis gravidarum is associated with increased risk of preterm pre-eclampsia, placental abruption and small for gestational age birth
- 75 Gestational diabetes: higher animal protein intake during pregnancy is associated with increased risk, and higher vegetable protein intake with decreased risk
- 76 Late pregnancy use of selective serotonin reuptake inhibitors and serotonin and norepinephrine reuptake inhibitors is associated with increased risk of postpartum haemorrhage
- 77 Antenatal perineal massage decreases risk of perineal trauma during birth
- 78 In Western Australia, 70% of mothers of babies with fetal alcohol syndrome did not have an alcohol-related diagnosis recorded during pregnancy
- 79 Adolescent girls who experience abuse or neglect are at an increased risk of teen pregnancy

### Mental health

- 80  People with schizophrenia are more likely to have a mother who smoked during pregnancy than people without the condition

### Adult nursing

- 81  Home palliative care services increase the chance of adults with advanced illness dying at home and reduce symptom burden without impact on caregiver grief

### Child health

- 82 Mandatory gloving in acute care paediatric units associated with decreased risk of hospital-acquired infections
- 83 Parents of British Bangladeshi children undergoing cancer treatment report managing competing knowledge, vigilance, advocacy, balancing work and treatment burden impacts on their lives

- 84 Prenatal exposure to sodium valproate is associated with increased risk of childhood autism and autistic spectrum disorder
- 85 Excess weight gain in the first 18 months of life is associated with later childhood overweight, obesity and greater arterial wall thickness
- 86 Children of mothers with iodine deficiency during pregnancy are more likely to have lower verbal IQ and reading scores at 8–9 years of age
- 87 Alcohol consumption in early and late pregnancy is associated with poor child reading and writing ability at 8–9 years of age
- 88 Children's experience of postoperative pain relief: children, parents and nurses use various pharmacological and non-pharmacological approaches, particularly distraction

### Care of people with learning disabilities

- 89 In adults with intellectual disability, discontinuation of antipsychotics is associated with reduction in weight, BMI, waist circumference and blood pressure

### Primary health care

- 90 Calcium supplements associated with increased risk of cardiovascular death in men but not women
- 91 One year following injury, pain and physical factors affect the likelihood of residual disability, but psychiatric symptoms may have a greater influence
- 92 Modest salt intake reduction for 4 weeks or longer decreases blood pressure

### Nursing issues


- 94 End-of-life care in the emergency department: nurses who invest in the nurse–patient relationship are better able to manage the emotional aspects of caring for dying people and their relatives
- 95 Terminally ill patients who are supported by religious communities are more likely to receive aggressive end-of-life care rather than hospice care; spiritual support from medical teams may reverse this
- 96 The majority of hospitalised elderly people at high risk of dying have thought about end-of-life care, though documentation of preferences in medical records may be lacking
- 97 Women's experience of abuse in a healthcare setting reveals preceding loss of power struggles, with sensitivity and dependency making them vulnerable to staff domination

### Care of the older person

- 98 A transitional care model for low-income older adults does not reduce readmission rates or emergency department visits during care transitions

### PostScript

- 66 Correction

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics  
[www.publicationethics.org.uk](http://www.publicationethics.org.uk)

 equator network

 recycle  
When you have finished with this please recycle it



Receive regular table of contents by email.  
Register using this QR code.