
Evidence-Based Nursing



Join Evidence-Based Nursing Twitter Journal Chat #ebnjc

Contents

Editorial

99 Legalising assisted suicide: what does the evidence say?

Research made simple

100 Bias in research



Women's health

102 Feasibility and efficacy of an internet treatment for postnatal depression utilising a behavioural activation approach

Midwifery

- 103 Women admitted to hospital during pregnancy have increased risk of venous thromboembolism that persists up to 28 days postdischarge
- 104 Pregnancy obesity is associated with increased rates of all-cause mortality and cardiovascular hospital admissions in adult offspring
- 105 Psychoeducation for pregnant women with fear of childbirth increases rates of spontaneous vaginal delivery, reduces caesarean rates and improves delivery experience
- Higher coffee intake in pregnancy linked to prolonged gestation, and higher caffeine intake linked with babies being small for gestational age

Adult nursing

- **107** Earlier thrombolytic treatment is associated with better outcomes following acute ischaemic stroke
- 108 People receiving dialysis in the morning have better subjective sleep quality than those who receive dialysis at other times
- 109 Review: exercise interventions improve pain and function in people with knee osteoarthritis compared with no exercise

Child health

- 110 Parental—adolescent conversations that focus on weight are more likely to be associated with unhealthy weight-control behaviours in adolescents than conversations that focus on healthy eating
- Parents of non-verbal children with learning disability (LD) most commonly recognise their child's pain through vocalisations, social behaviour and facial expressions
- 112 Overweight or obese young people are not at increased risk of depression, but young people with depression are at increased risk of obesity

October 2014 Vol 17 No 4

- Although parents are generally satisfied with their child's postoperative care, children continue to experience moderate-to-severe pain postoperatively
- 114 Child-spanking at age 5 may have a detrimental effect on child externalising behaviour and cognitive development at age 9
- 115 Infrared thermometers for assessing fever in children: the ThermoScan PRO 4000 ear thermometer is more reliable than the Temporal Scanner TAT-500

Health promotion

Few people living with and beyond cancer meet current exercise recommendations



Primary health care

117 A self-regulation protocol for people with COPD improves symptoms and reduces unscheduled physician visits

Nursing issues

- 118 Topical lidocaine—epinephrine—tetracaine is effective in reducing pain during laceration repair with tissue adhesive in children
- 119 No difference in patient satisfaction or quality of care provided by nurses or GPs for low-complexity primary care presentations
- 120 In menopausal women physical activity and taking dietary supplements may protect against coronary artery disease
- 121 Caring behaviour of nurses in Malaysia is influenced by spiritual and emotional intelligence, psychological ownership and burnout
- 122 Higher levels of nurse staffing are not associated with reduced adverse events among postoperative children if surveillance levels are low

Care of the older person

- 123 A model to develop compassionate relationship-centred care between older people, relatives and staff identifies seven factors necessary to promote appreciative caring conversations
- Around a quarter of all older people have disability during the last 2 years of life, with prevalence higher among women and those aged over 80 years
- 125 Exercise-based fall prevention programmes decrease fall-related injuries
- 126 Healthcare professionals may not be maintaining person-centred care for people with dementia hospitalised on acute wards



This article has been chosen by the Editor

to be of specialist interest or importance

and is freely available online



