


Evidence-Based Nursing



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc

Contents

Editorial

- 1 Changes to student nurse financing in the UK: a good or a bad thing?

Research made simple

- 2 Reviewing the literature



EBN opinion

- 4 Optimising palliative and end-of-life care within care home settings

Midwifery

- 5 Caesarean section and image consciousness among barriers to breast feeding for obese women
- 6 Greater support and debriefing may help student midwives to process traumatic birth experiences
- 7 Midwifery training needs identified when caring for women with female genital mutilation
- 8 Immediate preterm delivery for pregnancy hypertension increases risk of neonatal respiratory distress
- 9 Mothers of lower socioeconomic status make the decision to formula-feed in the context of culturally shared expectations and practices

Mental health

- 10 A body-mind-spirit intervention helps treat depression in India

Adult nursing

- 11 In people with atrial fibrillation receiving antithrombotics, short-term non-steroidal anti-inflammatory drug exposure increases risk of serious bleeding
- 12 Nurse-led home exercise programme improves physical function for people receiving haemodialysis

Child health

- 13 Parents are psychologically affected by their experiences when their child is in hospital because of uncertainty about prognosis and anxiety at the time of admission
- 14 Child health nurses miss opportunities to tackle obesity
- 15 Primary school education programmes can decrease child salt intake

January 2016 Vol 19 No 1

Care of people with learning disabilities

- 16 Guidelines improve nurses' knowledge and confidence in diagnosis of anxiety and challenging behaviours in people with intellectual disabilities

Health promotion

- 17 Smoking cessation strategies may increase quit rates among adults of low socioeconomic status
- 18 Daily e-cigarette use increases quit attempts and reduces smoking with no effect on cessation

Primary health care

- 19 In South Asian people with cardiac disease, strong patient-professional relationships may increase treatment compliance

Pain management

- 20 Nurses experience feelings of disempowerment when caring for patients in severe pain
- 21 No difference in pain, swelling or function with NSAIDs compared with paracetamol for soft tissue injury

Nursing issues

- 22 Productive hours by direct-care providers is the best measure of nurse staffing in relation to patient outcomes
- 23 Nursing students lack confidence in dealing with domestic abuse
- 24 Structured antibiotic 'time-out' audits as recommended by the CDC reduce antibiotic use and costs
- 25 Balance of lifetime remaining to anticipated suffering behind
- 26 Male victims of physical partner violence have poorer physical and mental health than men of the general population
- 27 After the Liverpool Care Pathway clear guidance and support on end-of-life care is needed

Care of the older person

- 28 Health professionals generally have more restrictive attitudes towards assisted dying in dementia than the public
- 29 Community palliative care use by dementia sufferers may reduce emergency department use at end of life
- 30 Increased age and understanding about treatment linked to better antihypertensive adherence
- 31 Non-pharmacological interventions for agitation in dementia: various strategies demonstrate effectiveness for care home residents; further research in home settings is needed
- 32 Training programmes and mealtime assistance may improve eating performance for elderly long-term care residents with dementia

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

C O P E COMMITTEE ON PUBLICATION ETHICS

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics
www.publicationethics.org.uk

 equator network

 recycle
When you have finished with this please recycle it



Receive regular table of contents by email.
Register using this QR code.