

Evidence-Based Nursing

July 2016 Vol 19 No 3

Contents

Editorial

- 65 Can the use of social media contribute towards Nursing and Midwifery Council revalidation?

Research made simple

- 66 Using observational research to obtain a picture of nursing practice



EBN opinion

- 68 Effective communicating starts with #hellomynameis

Women's health

- 70 Caesarean delivery is associated with higher risk of depressive symptoms, pain and sexual dysfunction
- 71 The beneficial effect of a Mediterranean diet supplemented with extra virgin olive oil in the primary prevention of breast cancer among women at high cardiovascular risk in the PREDIMED Trial

Midwifery

- 72 Continuity of midwifery care models improve outcomes for young women and babies
- 73 Organisation of postnatal care affects parents' confidence and security following hospital discharge at 72 h or less
- 74 Further research needed before late pregnancy ultrasound can be recommended for all

Adult nursing

- 75 Communication and support from healthcare professionals is essential for living kidney donors
- 76 Abdominal massage may decrease gastric residual volumes and abdominal circumference in critically ill patients
- 77 A modification to the Valsalva manoeuvre improves its effectiveness in treating supraventricular tachycardia
- 78 Fatigue 2 months after myocardial infarction may indicate risk for persistent fatigue
- 79 Psycho-educational programmes led by experienced nurses may improve the coping abilities of implantable cardioverter defibrillator recipients

Child health

- 81 Nasal balloon autoinflation can help clear middle ear effusion, improving the quality of life in school-aged children with glue ear

- 82 Legitimate opioid prescription increases the risk for future opioid misuse in some adolescents
- 83 Breast feeding could reduce the risk of childhood leukaemias

Primary health care

- 84 Health values change over time in people with type 2 diabetes
- 85 Combining nurse-led care with patient self-assessment could enhance tight control of rheumatoid arthritis and management of comorbidities

Pain management


- 87 Effects of an Internet-delivered cognitive behavioural therapy programme on chronic pain patients
- 88 Patients are aware of risks of opioid dependence, yet note poor communication from providers about pain and pain management

Nursing issues

- 89 Low-pressure, knee-length compression stockings provide most comfort with no effect on thrombosis risk
- 90 Emergency department nurses report high workload and management pressure to meet 4 h treatment targets
- 91 Good peer relationships can attenuate the negative effect of horizontal violence on job satisfaction
- 92 Patients' perspectives on the impact of fasting while in hospital
- 93 Costs incurred by family caregivers of patients with palliative care needs
- 94 Home death versus hospital death: are they comparable at all?

Care of the older person

- 95 Mealtime assistance may increase the energy and protein intake of hospitalised older patients
- 96 Implementation of a feasible monthly vitamin D intervention in homebound older adults using a Meals-on-Wheels programme



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

COPE

Member since 2010

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org.uk

equator network

recycle

When you have finished with this please recycle it



Receive regular table of contents by email.
Register using this QR code.