

Evidence-Based Nursing

July 2016 Vol 19 No 3

Contents

Editorial

- 65 Can the use of social media contribute towards Nursing and Midwifery Council revalidation?

Research made simple

- 66 Using observational research to obtain a picture of nursing practice



EBN opinion

- 68 Effective communicating starts with #hellomynameis

Women's health

- 70 Caesarean delivery is associated with higher risk of depressive symptoms, pain and sexual dysfunction
- 71 The beneficial effect of a Mediterranean diet supplemented with extra virgin olive oil in the primary prevention of breast cancer among women at high cardiovascular risk in the PREDIMED Trial

Midwifery

- 72 Continuity of midwifery care models improve outcomes for young women and babies
- 73 Organisation of postnatal care affects parents' confidence and security following hospital discharge at 72 h or less
- 74 Further research needed before late pregnancy ultrasound can be recommended for all

Adult nursing

- 75 Communication and support from healthcare professionals is essential for living kidney donors
- 76 Abdominal massage may decrease gastric residual volumes and abdominal circumference in critically ill patients
- 77 A modification to the Valsalva manoeuvre improves its effectiveness in treating supraventricular tachycardia
- 78 Fatigue 2 months after myocardial infarction may indicate risk for persistent fatigue
- 79 Psycho-educational programmes led by experienced nurses may improve the coping abilities of implantable cardioverter defibrillator recipients

Child health

- 81 Nasal balloon autoinflation can help clear middle ear effusion, improving the quality of life in school-aged children with glue ear

- 82 Legitimate opioid prescription increases the risk for future opioid misuse in some adolescents
- 83 Breast feeding could reduce the risk of childhood leukaemias

Primary health care

- 84 Health values change over time in people with type 2 diabetes
- 85 Combining nurse-led care with patient self-assessment could enhance tight control of rheumatoid arthritis and management of comorbidities

Pain management


- 87 Effects of an Internet-delivered cognitive behavioural therapy programme on chronic pain patients
- 88 Patients are aware of risks of opioid dependence, yet note poor communication from providers about pain and pain management

Nursing issues

- 89 Low-pressure, knee-length compression stockings provide most comfort with no effect on thrombosis risk
- 90 Emergency department nurses report high workload and management pressure to meet 4 h treatment targets
- 91 Good peer relationships can attenuate the negative effect of horizontal violence on job satisfaction
- 92 Patients' perspectives on the impact of fasting while in hospital
- 93 Costs incurred by family caregivers of patients with palliative care needs
- 94 Home death versus hospital death: are they comparable at all?

Care of the older person

- 95 Mealtime assistance may increase the energy and protein intake of hospitalised older patients
- 96 Implementation of a feasible monthly vitamin D intervention in homebound older adults using a Meals-on-Wheels programme



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc

This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org.uk



Receive regular table of contents by email.
Register using this QR code.