


# Evidence-Based Nursing



Join Evidence-Based Nursing  
Twitter Journal Chat #ebnjc

## Contents

### Editorial

- 97 Brexit: an uncertain journey

### Research made simple

- 99 Social media: the relevance for research



### Resources

- 101 Resources page

### EBN opinion

- 105 Relevance of social media to nurses and healthcare: 'to tweet or not to tweet'

### EBN perspectives

- 107 Perspectives on current child health issues

### Health promotion and public health

- 110 Parents who are aware when they are overweight themselves are more likely to recognise this in their children

### Women's health and midwifery

- 111 Postpartum depression can be prevented in low-income mothers using interpersonal psychotherapy
- 112 Oral fluconazole is often used during pregnancy in the treatment of vulvovaginal candidiasis despite safety concerns to the fetus
- 113 Trends in maternal morbidity relating to time of delivery need further exploration
- 114 'Time for self' appears to be a proactive strategy for the prevention of postpartum depression
- 115 Vaginal birth after caesarean at home in England: an opportunity for shared decision-making

### Mental health

- 116 Valuing therapeutic alliances with people experiencing acute psychosis

October 2016 Vol 19 No 4

### Adult nursing

- 117 Informal caregivers should be better supported—they are vital members of the healthcare team approach to chronic disease management

### Child health

- 118 Threats to parents' roles during the process of their child dying in the paediatric intensive care unit
- 119 Implementing preparatory strategies can help children participate in care
- 120 Group-based educational interventions in 11–16-year-olds can be effective in supporting suboptimal diabetes control

### Learning disabilities

- 122 Individualised approach to health promotion is needed for individuals with mild-to-moderate intellectual disability

### Nursing issues

- 123 Acceptance and commitment therapy: an appropriate treatment option for older adults with chronic pain
- 124 It is unclear if combined motivational interviewing and cognitive behavioural therapy improve medication adherence
- 125 Information gaps in medication communication during clinical handover calls for a different approach
- 126 Subclavian site should be preferred for central venous access

### Care of the older person

- 127 Hypovitaminosis D predicts more rapid and severe cognitive deterioration in ethnically diverse older adults with and without dementia
- 128 Nursing home staff should ensure that advance treatment decisions are discussed with residents and relatives

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

COPE

Member since 2010  
JMS06600

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics [www.publicationethics.org.uk](http://www.publicationethics.org.uk)

 equator  
network

 recycle  
When you have finished  
with this please recycle it



Receive regular table of contents by email.  
Register using this QR code.