

Evidence-Based Nursing

Contents

Purpose and procedure	98	Discharge planning and home follow up by advanced practice nurses reduced hospital readmissions of elderly patients	125
EBN notebook		Diagnosis	
The fundamentals of quantitative measurement	100	The Beck Depression Inventory for Primary Care accurately screened for major depressive disorders	126
Implementation forum		Prognosis	
Centres of evidence-based nursing: directions and challenges	102	Middle ear disease was associated with some behaviour problems at 5 and 10 years of age	127
Acknowledgments	104	Causation	
Other articles noted	105	A decrease in care was associated with suicide in people with mental illness	128
Treatment		Review: depression is associated with an increased risk of cardiovascular mortality and suicide	129
A structured discharge package given by a nurse reduced hospital readmission in children with asthma ..	113	Regular alcohol intake decreased risk of coronary heart disease events but not total mortality in men	130
Home based care improved glycaemic control and was cost effective in children with type 1 diabetes	114	Increased calcium intake was associated with decreased risk of ischaemic heart disease mortality in postmenopausal women	131
Coping skills training reduced haemoglobin A _{1c} and improved self efficacy in youths with diabetes	115	Qualitative	
A bereavement support group reduced grief and distress in bereaved, homosexual men	116	Parents of terminally ill adolescents experienced changed parental roles and a distorted time world	132
Lifestyle and structured exercise programmes increased physical activity and improved cardiorespiratory fitness	117	Men found the early months of first time fatherhood challenging	133
Group plus individual education improved quality of life and lung function in patients with asthma	118	A blend of comforting strategies and a form of team comforting were used during nasogastric tube insertion	134
Back pain self management groups led by lay people increased self care confidence and reduced activity limitations at 6 months	119	Patients with chronic obstructive pulmonary disease experienced ongoing challenges of preserving their personal integrity	135
Review: medication and cognitive behaviour therapy control symptoms of bulimia nervosa	120	Adjusting to the ambivalence of nursing home life was explained by the phrase <i>making the best of it</i>	136
A culturally and sex specific behavioural intervention reduced sexually transmitted diseases in minority women	121	Glossary	Inside back cover
Early oral feeding reduced recovery time after surgery for gynaecological malignancies	122		
A clinical pathway reduced length of stay, time to ambulation, and complications after hip and knee arthroplasty	123		
A home based intervention reduced hospital readmissions and mortality in patients with congestive heart failure	124		