


# Evidence-Based Nursing



Join Evidence-Based Nursing  
Twitter Journal Chat #ebnjc

## Contents

### Editorial

- 59 Time to take a stand on nurses' working conditions?

### EBN opinion

- 60 Mobilising identity through social media: psychosocial support for young people with life-limiting conditions

### Research made simple

- 63 Data collection in qualitative research



### EBN perspective

- 65 Nurse education

### Commentaries

#### Nursing issues

- 67 Medication errors may be reduced by double-checking method
- 68 Likelihood of nursing care being missed is influenced by several work-based factors
- 69 Higher vitamin D intake could improve sleep and cognitive outcomes in older adults with heart failure

#### Nurse education

- 70 Interventions for coping with bullying need further investigation and should be built into nursing curricula

#### Mental health

- 73 Risk of suicidal thinking in caregivers of people with dementia should be assessed and monitored by health professionals

#### Care of the older person

- 74 Prehospital physical function and strength can influence clinical outcomes of critically ill older adults

## July 2018 Volume 21 Issue 3

- 75 Exercise interventions reduce the risk of injurious falls among older adults

- 76 Inpatient mobilisation programmes targeting older patients are feasible, but further research is required to examine impact on patient outcomes

### Women's health & midwifery

- 77 Study suggests that pre-existing maternal epilepsy increases the risk of adverse perinatal outcomes
- 78 There remains a need for greater awareness and further research on fathers' stress in the perinatal period

### Adult nursing

- 79 Insufficient evidence to support ultrasonography as a method to rule out an improperly positioned nasogastric tube
- 80 Extended visitation policy may lower risk for delirium in the intensive care unit
- 81 Stroke self-management programmes could improve patient self-efficacy and satisfaction with self-management behaviours
- 82 Nurses should inform patients of the possibility of awareness during bronchoscopy performed with procedural sedation
- 84 Adolescents and young adults are a unique patient group whose diagnosis of cancer presents challenges that require in-depth, individual assessment
- 85 Nurses and physicians bring different perspectives to end-of-life decisions in intensive care units

### Learning disabilities

- 86 Adults with intellectual disabilities experience shame that leads to psychological distress and mental illness that requires assessment and treatment



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebn.bmj.com/site/about/guidelines.xhtml#open>

C O P E

Member since 2010  
JMK6666

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics [www.publicationethics.org.uk](http://www.publicationethics.org.uk)

equator  
network

recycle

When you have finished  
with this please recycle it