


Evidence-Based Nursing



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc

Contents

Editorials

- 87 Skills and simulation in nursing: a great opportunity or huge challenge?
- 89 Developing and sustaining a Community of Practice through Twitter for work-based learning

OPEN ACCESS

EBN Opinion

- 91 Engaging use of social media as a research tool to capture the daily life experiences of young people with chronic pain

EBN Perspective

- 94 Pain

Research made simple

- 96 Phenomenology as a healthcare research method
- 99 How to appraise quantitative research

EDITOR'S CHOICE

Resources page

- 102 Resources page

Commentaries

Care of the older person

- 104 Antihypertensive medications may reduce the risk of dementia in older African-Americans with hypertension

Women's health and midwifery

- 106 Fear of inducing guilt or shame affects midwives' communication about body weight with pregnant women

October 2018 Volume 21 Issue 4

Child health

- 107 Combining morphine and ibuprofen does not improve pain control compared with using either drug alone following musculoskeletal injury in children
- 109 Adolescent girls are susceptible to sexual pressure, coercion and victimisation and strategies need to be developed and aimed at males and females to prevent this
- 110 'Rooming-in' could be an effective non-pharmacological treatment for infants with neonatal abstinence syndrome

- 111 Effectiveness of therapeutic clowning on handwashing habits remains unknown

- 112 Non-pharmacological approaches to infantile colic: white noise versus swinging

Adult nursing

- 113 Fatigue is an important potential symptom of myocardial infarction in women
- 114 Professionals need to discuss more effectively the holistic needs of their patients regarding palliative care
- 115 Cervical spine can be assessed effectively by emergency department triage nurses

Nursing issues

- 116 There is a lack of preventive care and treatment interventions for urinary incontinence in hospitalised adults
- 117 Children's healthcare nurses should receive education in discussing intimate partner violence with new mothers
- 118 Coping methods used by emergency department nurses after a workplace violence experience could influence their intention to leave the hospital



This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.



This article has been made freely available online under the BMJ Journals Open Access scheme. See <http://ebn.bmj.com/site/about/guidelines.xhtml#open>



Member since 2010

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org.uk



When you have finished with this please recycle it