

Promoting patient involvement through person-centred handovers in nursing

10.1136/ebnurs-2018-102982



Lisa Kidd

Nursing & Healthcare, University of Glasgow School of Medicine Dentistry and Nursing, Glasgow, UK

Correspondence to: Dr Lisa Kidd, Nursing & Healthcare, University of Glasgow School of Medicine Dentistry and Nursing, Glasgow G12 8QQ, UK; Lisa.Kidd@glasgow.ac.uk

Commentary on: Kullberg A, Sharp L, Dahl O, *et al.* Nurse perceptions of person-centered handovers in the oncological inpatient setting - A qualitative study. *Int J Nurs Stud* 2018;86:44-51.

Implications for practice and research

- ▶ Person-centred handovers can enhance patient participation in healthcare decision-making, supporting the provision of high quality, safe person-centred care.
- ▶ Consideration is needed to how person-centred handovers are understood and implemented in clinical practice to ensure that they foster genuine participation and partnership working between nursing teams and patients.

Context

Nurse handovers are a routine form of communication and information exchange that occurs when one nurse hands over the responsibility of care for a patient to another nurse, for example at the end of a nursing shift.¹ In daily practice, different models of handover are used.¹ Evidence on nurses' perspectives of person-centred handover (PCH) models, which incorporate and promote a greater degree of patient involvement than traditional professional-professional handover models, however, is sparse.²

Methods

In Kullberg *et al.*'s² study, semi-structured interviews were used to explore nurses' (n=11) perceptions of the use and impact of PCH in two inpatient oncology wards in Sweden. In both wards, PCH were conducted at the patients' bedside at the change of the morning and evening shifts. The PCH began with introducing the patient, family members and the oncoming staff before inviting the patient to raise any concerns or questions. Once these were addressed, information about the patients' symptoms, concerns and their planned care was communicated to the group. The PCH aimed to avoid the use of medical jargon and offered repeated opportunities for patients and visitors to raise concerns or issues. Interviews were analysed using qualitative content analysis.

Findings

Overall, the nurses expressed positive experiences with the PCH. PCH was perceived as more efficient than the previous handover style,

helped nurses to get to know their patients quicker, strengthened team working and offered an opportunity for learning and teaching across nursing teams. Involving patients' perspectives in the handover process was viewed as valuable for enhancing patient safety and facilitated the provision of individualised care. Concerns about the PCH included; how participation was conceptualised by all members of the team, the extent to which the patient understood their involvement in the PCH, impact on workload and patient confidentiality.

Commentary

Patient and public involvement (PPI) in healthcare, service planning and health research has received significant attention over the past decade with proponents of PPI arguing that it enhances the quality and personalisation of service provision.³ Encouraging greater patient involvement in decision-making is seen as a marker of the provision of high quality and safe person-centred care.⁴⁻⁶ The challenges identified in Kullberg *et al.*'s study² resonate with the wider literature on patient participation and involvement where issues of uncertainty over what genuine 'participation' and partnership working looks like and how best to do this well in practice, were again highlighted.⁴⁻⁶ Power inequalities, highlighted in previous research,⁶ were also identified in Kullberg *et al.*'s study. Nurses perceived that the PCH, if conducted by staff who stood looking down on the patient in a bed, could serve to reinforce hierarchical structures and make patients feel inferior. Such findings emphasise the challenges involved in the implementation of 'involvement' initiatives, such as PCH, in practice to ensure that these do not become at risk of tokenism. Furthermore, genuine participation and partnership working can only be achieved if all 'partners' are comfortable and informed about the purpose and nature of their involvement. The findings from Kullberg *et al.*'s study provide a novel insight into nurses' perspectives of the use of PCH. Further research in this area to conceptualise what 'successful' patient participation in the handover process looks like, and how this can be best achieved, from both patients' and professionals' perspectives would make a valuable contribution to the evidence base on PCH.

Competing interests None declared.

Provenance and peer review Commissioned; internally peer reviewed.

© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

References

1. Smeulers M, Lucas C, Vermeulen H. Effectiveness of different nursing handover styles for ensuring continuity of information in hospitalised patients. *Cochrane Database Syst Rev* 2014;CD009979.
2. Kullberg A, Sharp L, Dahl O, *et al.* Nurse perceptions of person-centered handovers in the oncological inpatient setting: A qualitative study. *Int J Nurs Stud* 2018;86:44-51.
3. Øvretveit J. *Does improving quality save money? A review of the evidence of which improvements to quality reduce costs to health service providers.* London: Health Foundation, 2009.
4. Ocloo J, Garfield S, Dawson S, *et al.* Exploring the theory, barriers and enablers for patient and public involvement across health, social care and patient safety: a protocol for a systematic review of reviews. *BMJ Open* 2017;7:e018426.
5. Foot C, Gilbert H, Dunn P, *et al.* *People in control of their own health and care The state of involvement London:* Kinds Fund, 2014.
6. Ocloo J, Matthews R. From tokenism to empowerment: progressing patient and public involvement in healthcare improvement. *BMJ Qual Saf* 2016;25:626-32.