


Evidence-Based Nursing



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc

Contents

Editorial

- 93 Mental health in the time of COVID-19



OPEN ACCESS

Research made simple

- 94 What are the foundations of a good PhD?



EDITOR'S CHOICE

Resources

- 97 Resources page

Commentaries

Adult nursing

- 98 Paying attention to cultural context matters in advance care planning and palliative care
- 99 Multicomponent exercise programmes among very elderly hospitalised patients can be safe and effective in reversing functional decline

Mental health

- 100 The online treatment tool, 'Mood Mechanic', is a viable option for the treatment of anxiety and depression in young adults who do not undertake face-to-face treatment
- 101 Cognitive behavioural therapy is not effective for depression in advanced cancer but could help in anxiety or other psychological symptoms

Women's health and midwifery

- 102 There are both positive and negative implications to the use of mobile phones in the birth suite
- 103 Call for culturally sensitive comprehensive cancer control programmes in the Middle Eastern countries

Care of the older person

- 104 'Use it or lose it': the relationship between the frequency of word puzzle use and cognitive function
- 105 Comorbidity is a risk factor for poor quality of life in people with dementia
- 106 Place of death preference among people with dementia: an important conversation

Learning disabilities

- 107 One size does not fit all: mental health services access for people with intellectual/learning disability requires individualised planning and education/training programmes for health and social care providers

October 2020 Volume 23 Issue 4

Nursing issues

- 108 Timely psychological interventions can ameliorate symptoms of prolonged grief
- 109 Enhancing the role of nursing in primary care facilitates task sharing and addresses human resource shortages to achieve optimal population health outcomes
- 110 Higher prevalence of pressure ulcers in people receiving palliative care is not necessarily an indicator of poor care
- 111 More research is needed to understand how to provide optimal palliative dementia care to people living at home
- 112 Cigarette smoking increases persistent pain intensity and interference, impairs function and sleep
- 113 Walking may ameliorate fatigue in women receiving chemotherapy for breast cancer
- 114 Nurses should be empowered to be proactively involved in family decision meetings on intensive care unit patient care decisions

Nurse education

- 115 The problem of late dropout from nursing education is a serious one and educational simulation can be used as a solution
- 116 Nurse educators should use a range of approaches to teach students compassion
- 117 Nurturing cultural awareness needs to be explicitly addressed in undergraduate nurse curricula

Community and primary care nursing

- 118 Mindfulness training for primary care patients promotes chronic disease self-management behaviours

Health promotion and public health

- 119 Oral health should be considered when working with adults with intellectual disabilities, and larger, higher-quality studies in this area are needed
- 120 Are E-cigarettes more effective in supporting smoking cessation than nicotine-replacement therapy?

This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

This article has been made freely available online under the BMJ Journals open access scheme. See <http://ebn.bmj.com/site/about/guidelines.xhtml#open>



This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org.uk



When you have finished with this please recycle it