


Evidence-Based Nursing



Join Evidence-Based Nursing
Twitter Journal Chat #ebnjc

Contents

Editorial

- 29  Spotlight on maternal mental health: a prepandemic and postpandemic priority

OPEN ACCESS

Resources

- 31 Resources page

Research made simple

- 32  Five tips for developing useful literature summary tables for writing review articles

EDITOR'S CHOICE

Commentaries

Child health

- 35 Association between COVID-19 and Kawasaki-like disease in children is a topic that needs further investigation

- 36 Direct-to-consumer telemedicine visits demonstrate decreased antibiotic prescribing quality in paediatric clients with acute respiratory infections

Health promotion and public health

- 37 Increased sedentary behaviour is associated with increased depression symptoms in adults

- 38 Evidence from randomised trials linking intake of red meat to diseases—including cardiovascular disease and cancer—is weak

- 39 Reducing consumption of red meat may have a very small effect on all-cause mortality and cardiometabolic diseases

Nurse education

- 40 Assertiveness training during early undergraduate years has the potential to increase students' confidence and ability to act as patient advocates

Nursing issues

- 41 Use of personal protective equipment reduces the risk of contamination by highly infectious diseases such as COVID-19

- 42 Complex interactions between provider, organisation and patient level characteristics influence the effect of pressure ulcer preventive care interventions

- 43 A workplace sales ban and motivational intervention can reduce sugar-sweetened beverage intake and improve cardiometabolic health

- 44 Patients need more than just verbal instructions upon discharge from the emergency department

- 45  Without adequate nursing support for families, dying at home threatens the values of a good death

EDITOR'S CHOICE

- 46  Social media is a source of health-related misinformation

EDITOR'S CHOICE

- 47 Better patient activation is a precursor to engagement in shared decision making

April 2021 Volume 24 Issue 2

Learning disabilities

- 48 Time to increase the involvement of persons with intellectual disability in palliative care and research

- 49 Ethical dilemmas and legal ambivalence affects the midwives' perception for providing sexual and reproductive health consultation to women with intellectual disability

Care of the older person

- 50  Feeling fearful and lonely are indicative experiences of emotional distress for people with dementia

OPEN ACCESS

- 52 Family caregivers expect a patient and family centric approach in holistic end-of-life provision

- 53  Hopelessness during acute hospitalisation is a strong predictor of mortality

OPEN ACCESS

- 54 Delirium occurrence in hospitalised older people is positively associated with development of dementia postdischarge

- 55 Relational care in older persons nursing is not just a consideration but an organisational priority

- 56 Frailty is associated with a higher use of potentially inappropriate medications among community-dwelling older adults

Women's health and midwifery

- 57 Uncovering the use and misuse of opioid analgesia in a pregnant and non-pregnant female population

- 58 Cannabis use in pregnancy: a harm reduction approach is needed with a focus on prevention and positive intervention

- 59 COVID-19 and pregnancy outcomes: initial findings show little threat, but more data are needed

Adult nursing

- 60 Survivorship care plans have minimal impact on health actions

- 61 Current state of knowledge around nurse-led cardioversion: a launchpad for future projects

- 62 Shorter intervals and sequential administration of adjuvant chemotherapy are effective in reducing the 10-year risk of recurrence and death in early breast cancer women without detrimental effects

- 64 Nurses' intuitive worrying predicts physiological changes in patients' vital signs

- 65 Bariatric surgery could reduce the risk of specific types of breast cancer

Mental health

- 66 Peer mentorship of hospital inpatients with substance use disorders is associated with beneficial outcomes

- 67 Short therapy programmes may reduce the risk of suicide reattempts

- 68 Depressive symptoms in the last year of life: early screening and varied treatment pathways needed

 This article has been chosen by the Editor to be of specialist interest or importance and is freely available online.

 This article has been made freely available online under the BMJ Journals open access scheme. See <http://ebn.bmj.com/site/about/guidelines.xhtml#open>

C O P E

Member since 2010
JMK06600

This journal is a member of and subscribes to the principles of the Committee on Publication Ethics www.publicationethics.org

 **recycle**
When you have finished
with this please recycle it