

Evidence-Based Nursing

Contents

| | |
|--|-------------------|
| Purpose and procedure | 34 |
| EBN notebook | |
| Clinically useful measures of the effects of treatment | 36 |
| Sigma Theta Tau contest | |
| Evidence-based practice contest: innovations in clinical excellence | 40 |
| Treatment | |
| Routine primary immunisation using a longer needle resulted in fewer local reactions in infants | 41 |
| Few young children had mild reactions to the measles, mumps, and rubella vaccine and older children had nearly no vaccine related reactions | 42 |
| Review: recall or reminder systems improve the rate of childhood and adult immunisation | 43 |
| A clinical pathway reduced use of nebulised β agonists and length of hospital stay in children with asthma exacerbations | 44 |
| Review: brief primary care interventions are moderately effective for increasing physical activity | 45 |
| Contamination of urine specimens from women with acute dysuria did not differ with collection technique | 46 |
| Review: NSAIDs, paracetamol, and analgesic/anti-inflammatory oral rinse reduce sore throat symptoms .. | 47 |
| Docusate sodium with irrigation was better than triethanolamine polypeptide with irrigation for dissolving earwax | 48 |
| Early feeding after major non-laparoscopic gynaecological surgery did not increase gastrointestinal symptoms | 49 |
| Adjunctive nurse telehealth care reduced depressive symptoms and improved functioning | 50 |
| A shared care system of hospital follow up reduced pain and use of healthcare resources and increased satisfaction in patients with rheumatoid arthritis | 51 |
| A cognitive behavioural family intervention reduced psychiatric morbidity in caregivers of patients with Alzheimer's disease | 52 |
| Integrated care pathway based rehabilitation for acute stroke did not reduce length of hospital stay | 53 |
| Intensive geriatric rehabilitation reduced hospital stay and time to independent living in hip fracture patients with mild to moderate dementia | 54 |
| Assessment | |
| A clear urine specimen on visual inspection cannot totally exclude a diagnosis of urinary tract infection | 55 |
| Causation | |
| Children who were exposed to antenatal corticosteroids and had birth weights ≤ 1500 g did not have adverse growth, cognitive, or lung function outcomes at age 14 years | 56 |
| Quality improvement | |
| A nurse led clinic and computer decision support system for anticoagulation decisions was at least as effective as a hospital clinic | 57 |
| Clinical prediction guide | |
| An antenatal index predicted postpartum depression with high sensitivity and moderate specificity | 58 |
| Qualitative | |
| Late adolescent girls used 4 stories to describe their smoking patterns | 59 |
| Teenage mothers with oppressive pasts created new family caregiving traditions | 60 |
| Individuals with HIV/AIDS ascribed several different meanings to their use of complementary therapy | 61 |
| Inpatients had mostly negative experiences of seclusion during short term treatment in a mental health facility | 62 |
| Properly planned de-institutionalisation for mental illness maintained most in community living with enhanced quality of life | 63 |
| Elderly women described 5 phases of living with change after myocardial infarction | 64 |
| Glossary | Inside back cover |