Prognosis

Needing help to use a toilet was associated with faecal incontinence at 3 months in new onset faecal incontinence after stroke

Harari D, Coshall C, Rudd, AG, et al. *New-onset fecal incontinence after stroke: prevalence, natural history, risk factors, and impact.* *Stroke* 2003;34:144–50.

QUESTION: In patients with new onset faecal incontinence after stroke, which prognostic factors are associated with bowel continence status at 3 months of follow up?

Design

Inception cohort followed up for 3 years.

Setting

South London, UK.

Patients

1468 patients without pre-existing faecal incontinence, who survived a first in a lifetime stroke were recruited from the community based South London Stroke Register. 846 patients (53% men) were included in the primary analysis at 3 months (412 patients had died, and data for 210 patients were not available).

Assessment of prognostic factors

Data were collected in the acute stroke phase (within the first 10 d) by research registrars, and at 3 months, 1 year, and then annually by face to face interviews with a trained field worker. When reliable answers could not be obtained directly from patients (because of cognitive impairment, communication difficulties, or severe illness), the information was sought from family, caregivers, nursing home or hospital staff, or community health workers. Medical information was also obtained from hospital and general practitioner records. Associations between faecal incontinence and prognostic factors were assessed in a cross sectional analysis using bivariate and multivariate statistical methods.

Main outcome measure

Bowel continence status measured by the Barthel Index bowel subscore at 7–10 days, 3 months, and 1, 2, and 3 years.

Main results

Prevalence of faecal incontinence after stroke was 30% at 7–10 days, 11% at 3 months and 1 year, and 15% at 3 years. 35% of patients with faecal incontinence at 3 months were continent by 1 year; 63% of patients with faecal incontinence at 1 year had been continent at 3 months. Of patients who were continent at 3 months, 6% became incontinent at 1 year.

Multivariate analysis of acute stroke characteristics showed that initial urinary incontinence (odds ratio [OR] 6.2, 95% CI 3.2 to 11.9) and visual and/or sensory neglect (OR 1.9, CI 1.0 to 3.5) were associated with faecal incontinence. After further adjustment for significant clinical and functional variables, needing help to use a toilet (OR 3.5, CI 1.4 to 17.3) and use of anticholinergic drugs (OR 3.1, CI 1.1 to 10.2) were independent prognostic factors for faecal incontinence.

Conclusion

New onset faecal incontinence in stroke survivors was common but may be transient. Needing help to use a toilet and use of anticholinergic drugs were associated with new onset faecal incontinence at 3 months after stroke.

COMMENTARY

The study by Harari *et al* enhances previous research on faecal incontinence in patients with stroke as it includes only patients with newly acquired faecal incontinence after stroke. Most importantly for healthcare professionals, this study assessed the contribution of potentially modifiable risk factors such as functional difficulties with toilet access and use of drugs that could lead to constipation with overflow.

Bowel continence status was assessed using a question from the Barthel Index. Patients were classified into 1 of 3 categories: “never” losing bowel control, “occasional accident,” or “all the time.” Although the validity of the Barthel Index as a functional measure has been shown, it may be limited in its ability to assess faecal incontinence. This limitation is apparent when one considers that a patient may have had only 1 faecal incontinence episode and yet be classified within the faecal incontinent sample. This classification may increase the similarity between the 2 groups (never experiencing faecal incontinence *v* experiencing faecal incontinence), and thus, any true association would be underestimated. A more sensitive measure of bowel continence would have strengthened this study. Furthermore, the lack of precision within the confidence intervals around the odds ratios for anticholinergic drug use and needing help with toilet use reflects substantial variability of the results. Thus, these results should be regarded as preliminary.

The results of this study are relevant to nurses, nurse practitioners, and clinical nurse specialists working with patients on stroke rehabilitation units and complex continuing care or long term care facilities. Because >10% of patients with stroke will have faecal incontinence, it is important to identify those individuals at greatest risk. In particular, the results show that faecal incontinence may be transient and modifiable. This will assist nurses to develop care plans to strengthen functional ability, particularly with respect to toileting. The findings also emphasise the importance of assessing modifiable risk factors in addition to clinical and demographic factors.

Nancy J Martin, RN, PhD
Director, Research and Performance Metrics
Grand River Hospital
Kitchener, Ontario, Canada

Sources of funding:
Action Research,
Northern and Yorkshire
Research and
Development
Cardiovascular
Diseases Programme,
Charitable Foundation
of Guy's and St
Thomas' Hospital, and
Stanley Thomas
Johnson Foundation.

For correspondence:
Dr D Harari,
Department of
Healthcare of the
Elderly, Guy's and St
Thomas' Hospitals
Trust, London, UK.
danielle.harari@kcl.ac.uk